

Lecture PowerPoint

Chapter 27

Physics: Principles with Applications, 6th edition


Giancoli

© 2005 Pearson Prentice Hall

This work is protected by United States copyright laws and is provided solely for the use of instructors in teaching their courses and assessing student learning. Dissemination or sale of any part of this work (including on the World Wide Web) will destroy the integrity of the work and is not permitted. The work and materials from it should never be made available to students except by instructors using the accompanying text in their classes. All recipients of this work are expected to abide by these restrictions and to honor the intended pedagogical purposes and the needs of other instructors who rely on these materials.

Chapter 27

Early Quantum Theory and Models of the Atom



Units of Chapter 27


Origins of the Quantum theory.....

- **Discovery and Properties of the Electron**
- **Planck's Quantum Hypothesis; Blackbody Radiation**
- **Photon Theory of Light and the Photoelectric Effect**
- **Energy, Mass, and Momentum of a Photon**
- **Photon Interactions; Pair Production**
- **Wave Nature of Matter**
- **Electron Microscopes**
- **Early Models of the Atom**
- **Atomic Spectra: Key to the Structure of the Atom**
- **The Bohr Model**
- **de Broglie's Hypothesis Applied to Atoms**

27.1 Discovery and Properties of the Electron


In the late 19th century, discharge tubes were made that emitted “cathode rays.”

Cathode rays are called electrons.



27.1 J.J Thomson's 1890 apparatus to measure the charge/mass ratio (e/m)

- Cathode rays could be deflected by electric or magnetic fields.
- Measuring the angle of deflection gave e/m
- But finding e and m separately proved elusive



27.1 Discovery and Properties of the Electron

By accelerating cathode rays through a known potential and then measuring the radius of their curved path in a known magnetic field, the charge to mass ratio could be measured:

$$v = \frac{E}{B}$$
$$\frac{e}{m} = \frac{E}{B^2 r} \quad (27-1)$$


The result is $e/m = 1.76 \times 10^{11} \text{ C/kg}$

Cathode rays are called electrons.

27.1 Discovery and Properties of the Electron

Millikan devised an experiment to measure the charge on the electron by measuring the electric field needed to suspend an oil droplet of known mass between parallel plates.

The mass and charge of each oil droplet were measured; careful analysis of the data showed that the charge was always an integral multiple of a smallest charge, e .



$$e = 1.602 \times 10^{-19} \text{ C}$$

Knowing e allows the electron mass to be calculated:

$$m_e = 9.11 \times 10^{-31} \text{ kg}$$

Design of the e/m experiment



- Accelerate a beam of electrons through a known potential difference so they gain a known kinetic energy ($E = qV$) and speed.
- Direct the beam through a known magnetic field, so the beam bends into a circle
- Measure the radius of the circular path
- Equate the centripetal force to the magnetic force

Voltage to Accelerate electrons


Current to coils, to provide magnetic field

27.2 Planck's Quantum Hypothesis; Blackbody Radiation

Blackbody radiation curves for three different temperatures.

Wien's Law:

$$\lambda_p T = k = 2.9 \times 10^{-3} \text{ m.K}$$




Copyright © 2005 Pearson Prentice Hall, Inc.

Radiation and Temperature

Heated bodies generally radiate across the entire electromagnetic spectrum

There is one particular wavelength, λ_p , at which the radiation is most intense.



27.2 Planck's Quantum Hypothesis; Blackbody Radiation

This spectrum could not be reproduced using 19th-century physics.

All attempts were either purely descriptive (like Wien's law), or led to the so called "ultraviolet catastrophe" - failure to derive the decline in flux at short wavelengths, leading to unrealistic prediction of limitless radiation in the UV.

A solution was proposed by Max Planck in 1900:

The atoms are all radiating, absorbing and redistributing energy between themselves. Each behaves as a harmonic oscillator with discrete modes

The distribution of atomic oscillator energies leads to the black-body spectrum

The oscillations within atoms can only occur at discrete frequencies that are multiples of a certain minimum value. (like modes on a string, or in a pipe)

Introduced a new fundamental constant of nature: Planck's Constant: h

Many including Plank viewed this model as purely a mathematical device

Today we view it as the first physically correct model in quantum physics!

27.2 Planck's Quantum Hypothesis; Blackbody Radiation

Planck found the value of his constant by fitting blackbody curves: $h = 6.626 \times 10^{-34} \text{ J}\cdot\text{s}$

Planck's proposal was that the energy of an oscillation had to be an integer multiple of hf . This is called the quantization of energy.

$$E = hf$$

$$E = nhf \quad \text{with } n = 1, 2, 3, \dots$$

27.3 Photon Theory of Light and the Photoelectric Effect

Einstein (1905) suggested that, given the success of Planck's theory, light must be emitted in small energy packets:

$$E = hf \quad (27-4)$$

These tiny packets, or particles, are called photons.

This theory won Einstein the Nobel Prize

He had resurrected Newton's 400 yr old idea of light as a stream of particles

27.3 Photon Theory of Light and the Photoelectric Effect


The photoelectric effect:

If light strikes a metal, electrons are emitted.

$$hf = KE_{max} + W_0$$

$$KE_{max} = eV_0$$

The effect does not occur if the frequency of the light is too low



27.3 Photon Theory of Light and the Photoelectric Effect


If light is a stream of particles, one predicts:

- Increasing intensity increases number of electrons but not energy**
- Above a minimum energy required to break atomic bond, kinetic energy will increase linearly with frequency**
- There is a cutoff frequency below which no electrons will be emitted, regardless of intensity**

27.3 Photon Theory of Light and the Photoelectric Effect

The particle theory assumes that an electron absorbs a single photon.


Plotting the kinetic energy vs. frequency:



This shows clear agreement with the photon theory, and not with wave theory.

27.3 Photon Theory of Light and the Photoelectric Effect

The photoelectric effect is how “electric eye” detectors work. It is also used for movie film soundtracks.



Copyright © 2005 Pearson Prentice Hall, Inc.

Photoelectric Effect also occurs in Nature -e.g. Photosynthesis, Sunburn

27.4 Energy, Mass, and Momentum of a Photon

- A photon must travel at the speed of light.
- Quantum theory (photoelectric effect) says
Kinetic energy = $E_{\text{photon}} = hf$
- Look at the relativistic equation for momentum:
 $E^2 = p^2c^2 + m^2c^4$
- The photon has no mass, so this reduces to
 $E_{\text{photon}} = pc$
- We combine these equations and find that a photon must carry momentum!

$$p = \frac{E}{c} = \frac{hf}{c} = \frac{h}{\lambda} \quad (27-6)$$

Summary of Chapter 27

- **Planck's hypothesis: molecular oscillation energies are quantized**

$$E = nhf$$

- **Light can be considered to consist of photons, each of energy**

$$E = hf$$

- **Photoelectric effect: incident photons knock electrons out of material**

- **Compton effect and pair production also support photon theory**

- **Wave-particle duality – both light and matter have both wave and particle properties**


- **Wavelength of an object:** $\lambda = \frac{h}{p}$

27.10 Early Models of the Atom

It was known that atoms were electrically neutral, but that they could become charged, implying that there were positive and negative charges and that some of them could be removed.

← $\approx 10^{-10}$ m →

One popular atomic model was the “plum-pudding” model:





Positively
charged
material

Rutherford's Experiment


- Rutherford in 1911 did an experiment that showed that the positively charged nucleus must be extremely small compared to the rest of the atom, and contain essentially all of the mass.
- He directed alpha particles (emitted by some radioactive elements) at a thin metal foil and observed the scattering angle.
- Rutherford found that a tiny fraction of these “bullets” bounced right back!
- Meaning they had hit something massive, yet tiny.
- the atom was mostly empty space!

Rutherford's Experiment

The only way to account for the large angles was to assume that all the positive charge was contained within a tiny volume – now we know that the radius of the nucleus is 1/10000 that of the atom



Copyright © 2005 Pearson Prentice Hall, Inc.



(b)

Copyright © 2005 Pearson Prentice Hall, Inc.

• **A BIG problem: Accelerated charges radiate EM radiation**


• **if the electron loses potential energy it falls closer to the nucleus.....**

• **So this electron would spiral into the nucleus immediately**


27.11 Atomic Spectra: Key to the Structure of the Atom

An atomic spectrum is a line spectrum – only certain frequencies appear.

If white light passes through such a gas, it absorbs at those same frequencies.



(a) **Hydrogen**



(b) **Helium**





7 \ **The Sun**

Each line is due to photons with a specific energy

What if these energies correspond to special “allowed” level inside the atom?

27.11 Atomic Spectra: Key to the Structure of the Atom

A very thin gas heated in a discharge tube emits light only at characteristic frequencies.



27.11 Atomic Spectra: Key to the Structure of the Atom

The wavelengths of electrons emitted from hydrogen have a regular pattern:

$$\frac{1}{\lambda} = R \left(\frac{1}{2^2} - \frac{1}{n^2} \right), \quad n = 3, 4, \dots \quad (27-9)$$

This is called the Balmer series. R is called the Rydberg constant:

$$R = 1.0974 \times 10^7 \text{ m}^{-1}$$

- **A spectral lines occurs for each value of n .**
- There are 5 visible lines in the Balmer series: $H\alpha$ (red), $H\beta$, $H\gamma$, $H\delta$, $H\epsilon$
- This is a ‘phenomenological’ or ‘empirical’ theory -it is very accurate but explains nothing. The constant “ R ” is easily measured with a spectroscope (basically a prism or a diffraction grating)
- But we are still at “WHY?”

27.11 Atomic Spectra: Key to the Structure of the Atom

Other series include the Lyman series (in the Ultraviolet):

$$\frac{1}{\lambda} = R \left(\frac{1}{1^2} - \frac{1}{n^2} \right), \quad n = 2, 3, \dots$$


And the Paschen series (in the Infrared):

$$\frac{1}{\lambda} = R \left(\frac{1}{3^2} - \frac{1}{n^2} \right), \quad n = 4, 5, \dots$$

Note the series differ only in this number


27.11 Atomic Spectra: Key to the Structure of the Atom

A portion of the complete spectrum of hydrogen is shown here. The lines cannot be explained by the Rutherford theory.



27.12 The Bohr Atom

Niels Bohr proposed that the possible energy states for atomic electrons were quantized – only certain values were possible. Then the spectrum could be explained as transitions from one level to another.



27.12 The Bohr Atom

After trying various ideas....

Bohr found that the angular momentum was quantized:


$$L = mvr_n = n \frac{h}{2\pi}, \quad n = 1, 2, 3, \dots \quad (27-11)$$

It worked perfectly!

But there was still no physical motivation.....

27.12 The Bohr Atom

An electron is held in orbit by the Coulomb force:



So you can calculate the radii of the “orbit” responsible for each spectral line.

Copyright © 2005 Pearson Prentice Hall, Inc.

By equating centripetal force and coulomb force

27.12 The Bohr Atom

Using the Coulomb force, we can calculate the radius of 1st orbit (n=1) orbit for hydrogen (Z=1):

$$r_1 = \frac{h^2}{4\pi^2 m k e^2} = 0.529 \times 10^{-10} \text{ m} \quad (27-13)$$

Radius of nth orbit:

$$r_n = \frac{n^2 h^2}{4\pi^2 m k Z e^2} = \frac{n^2}{Z} r_1 \quad (27-12)$$

The equation above comes from equating the centripetal force needed to balance the coulomb force. -a type of calculation you are all familiar with.

The resulting levels are not evenly spaced in radius- just like the spectral lines are not evenly spaced, but follow a distinct pattern

27.12 The Bohr Atom

For Hydrogen (Z=1)

- Lowest energy (n=1) → ground state energy:


$$E_1 = -13.6 \text{ eV}$$

- Higher energy states → excited states:

$$E_n = \frac{-13.6 \text{ eV}}{n^2} \quad n = 2, 3, 4 \dots$$

- Ionization energy or binding energy → Minimum energy required to remove an electron from the ground state
 - Requires +13.6eV to remove an electron from the ground state (n=1) for hydrogen.

27.12 The Bohr Atom



- The lowest energy level is called the ground state; the others are excited states.

- Notice how the levels are not evenly spaced in energy or in radius.

$$\frac{1}{\lambda} = R \left(\frac{1}{2^2} - \frac{1}{n^2} \right), \quad n = 3, 4, \dots \quad (27-9)$$

$$R = 1.0974 \times 10^7 \text{ m}^{-1}$$

Summary of Chapter 27

- Planck's hypothesis: molecular oscillation energies are quantized

$$E = nhf \quad n = 1, 2, 3, \dots$$

- Light can be considered to consist of photons, each of energy $E = hf$
- Photoelectric effect: incident photons knock electrons out of material
- Compton effect and pair production also support photon theory
- Wave-particle duality – both light and matter have both wave and particle properties
- Wavelength of an object: $\lambda = \frac{h}{p}$
- Principle of complementarity: both wave and particle properties are necessary for complete understanding
- Rutherford showed that atom has tiny nucleus
- Line spectra are explained by electrons having only certain specific orbits

Summary of Chapter 27

- Ground state ($n = 1$) has the lowest energy of -13.6 eV ; the others are called excited states. For Hydrogen ($Z = 1$)

$$E_n = \frac{-13.6 \text{ eV}}{n^2} \quad n = 2, 3, 4 \dots$$

- Wavelength emitted for hydrogen:

- Balmer series (in the visible) $\frac{1}{\lambda} = R \left(\frac{1}{2^2} - \frac{1}{n^2} \right), \quad n = 3, 4, \dots$

- Lyman series (in the ultraviolet) $\frac{1}{\lambda} = R \left(\frac{1}{1^2} - \frac{1}{n^2} \right), \quad n = 2, 3, \dots$

- Paschen series (in the infrared) $\frac{1}{\lambda} = R \left(\frac{1}{3^2} - \frac{1}{n^2} \right), \quad n = 4, 5, \dots$

- Rydberg constant: $R = 1.0974 \times 10^7 \text{ m}^{-1}$

ConceptTest 27.1

Photons

Which has more energy, a
photon of:

- 1) red light
- 2) yellow light
- 3) green light
- 4) blue light
- 5) all have the same energy



400 nm

500 nm

600 nm

700 nm

ConceptTest 27.1

Photons

Which has more energy, a photon of:

$$E = hf$$

- 1) red light
- 2) yellow light
- 3) green light
- 4) blue light
- 5) all have the same energy



400 nm

500 nm

600 nm

700 nm

The photon with the **highest frequency** has the **most energy** because $E = hf = hc/\lambda$ (recall that $c = f\lambda$). So a **higher frequency** corresponds to a **lower wavelength**. The highest energy of the above choices is **blue**.