

Chemical Oceanography Organics III

Dr. David K. Ryan


Department of Chemistry


University of Massachusetts Lowell

&

Intercampus Marine Science (IMS) Program










http://faculty.uml.edu/David_Ryan/84.653






Humification of
Organic Matter
(another scheme)

Libes, 1992


Group	Structure	pK _a	Hydrolysis Products	Exchange H ?
Alcohol		12	None	Yes
Phenol		10	None	Yes
Ether			None	
Aldehyde			None	No
Ketone			None	
Carboxyl		5	None	Yes
Ester			Carboxyl + Alcohol	
Amine		10	None	Yes
Amide			Carboxyl + Amine	Yes

Emerson & Hedges
Figure 8.2




Morel & Hering,
1993
See also
Emerson & Hedges
Figure 8.11 & 8.12

Figure 6.13 A possible pathway for the formation of marine humic acids from a triglyceride. From Harvey et al., 1983.




Humic Structure Proposed by Schnitzer (Rashid 1985)





Structure Attributed to Gamble et al. (1985)


Four possible carbon skeletons for the planar fulvic acid molecule




Morel & Hering (1993) Based on Aiken et al. (1985)




Possible Structural Units Set Forth by Averett, Leenheer, McKnight & Thorn (1989) From Morel & Hering, 1993





Organic Solute
Macromolecule
(ORMAC)
Leenheer 1985)

Molecular model of the lowest energy conformation of humic acid building blocks



Carbon atoms-green
Oxygen atoms-red
Nitrogen-blue
Hydrogen not shown

Davies & Ghabbour, 1999

Importance of Humic Materials

Global Carbon Reservoir

Take Part in Interfacial Phenomena

Undergo Coagulation and Aggregation

Involved in Photochemical Reactions

Contain Radicals

Known Reducing Agents

Methylate Metals

Form Chlorinated Species, THMs DBPs

Detoxify Metals

Limit Bioavailability of Metals

Alter Solubility

Influence

Bind Metals & Organic Pollutants


Terminal Electron Transport Acceptor for Bacteria



Sea Foam
caused by
naturally
occurring
surface
active
agents



Air-Sea Interfacial Chemistry



Importance of Humic Materials

Global Carbon Reservoir

Take Part in Interfacial Phenomena

Undergo Coagulation and Aggregation

Involved in Photochemical Reactions

Contain Radicals

Known Reducing Agents

Methylate Metals

Form Chlorinated Species, THMs DBPs

Detoxify Metals


Limit Bioavailability of Metals

Alter Solubility

Influence

Bind Metals & Organic Pollutants

Terminal Electron Transport Acceptor for Bacteria



Fox, 1983

Figure 3. Humic acid carbon as a function of salinity. The standard error of the mean is 5%.

Importance of Humic Materials

Global Carbon Reservoir

Take Part in Interfacial Phenomena

Undergo Coagulation and Aggregation

Involved in Photochemical Reactions

Contain Radicals

Known Reducing Agents

Methylate Metals

Form Chlorinated Species, THMs DBPs

Detoxify Metals

Limit Bioavailability of Metals

Alter Solubility

Influence

Bind Metals & Organic Pollutants

Terminal Electron Transport Acceptor for Bacteria

TABLE 10.2**Photoreactions of Organic Compounds**

Chromophore	Products or effects
Humic, fulvic	<ol style="list-style-type: none">1. Bleaching of absorption and fluorescence2. Production of singlet oxygen3. Fe(III) reduction4. Release of soluble P5. Oxidation of cumene via ROO and OH radicals6. Oxidation of phenolic groups to ArO and formation of e^- and O_2^-7. CO formation8. H_2O_2 formation (via O_2^- ?)
Chlorophyll	Loss of chlorophyll
Vitamins	Loss of bioassay activity
Amino acids	?
Glycine	COOH C-14 loss, HCHO 1 formation
$CH_3SSCH_3CH_3S$	CH_3S
CH_3ICH_3	CH_3
Fatty acids	Particles, absorb., hydroperoxides
Aldehydes	RCO, R, CO

Millero, 1996

Photochemistry

**CDOM = Chromophoric (or Colored)
Dissolved Organic Matter**




Figure 1. Absorption spectra (pathlength of 1 cm) of several waters and a generalized surface solar energy distribution (adapted from ref. 8). (DOC of waters: A = 3.0, B = 7.8, C = 13.4, D = 13.4, E = 15.4 mg L^{-1}).

Importance of Humic Materials

Global Carbon Reservoir

Take Part in Interfacial Phenomena

Undergo Coagulation and Aggregation

Involved in Photochemical Reactions

Contain Radicals

Known Reducing Agents

Methylate Metals

Form Chlorinated Species, THMs DBPs

Detoxify Metals

Limit Bioavailability of Metals


Alter Solubility

Influence

Bind Metals & Organic Pollutants

Terminal Electron Transport Acceptor for Bacteria

Quinone radical present in humic material



benzoquinone

semiquinone

hydroquinone

Scott, McKnight, Blunt-Harris, Kolesar & Lovely (1998) Environ. Sci. Technol. 32, 19

Importance of Humic Materials

Global Carbon Reservoir

Take Part in Interfacial Phenomena

Undergo Coagulation and Aggregation

Involved in Photochemical Reactions

Contain Radicals

Known Reducing Agents

Methylate Metals

Form Chlorinated Species, THMs DBPs

Detoxify Metals

Limit Bioavailability of Metals

Alter Solubility

Influence Transport

Bind Metals & Organic Pollutants

Terminal Electron Acceptor for Bacteria

Humics involved in many reduction reactions

- # Cr(IV) to Cr(III)
- # Fe(III) to Fe(II)
- # Hg(II) to Hg⁰
- # As, Se and V species

Importance of Humic Materials

Global Carbon Reservoir

Take Part in Interfacial Phenomena

Undergo Coagulation and Aggregation

Involved in Photochemical Reactions

Contain Radicals

Known Reducing Agents

Methylate Metals

Form Chlorinated Species, THMs DBPs

Detoxify Metals

Limit Bioavailability of Metals


Alter Solubility

Influence Transport


Bind Metals & Organic Pollutants

Terminal Electron Acceptor for Bacteria

Metal Complexation by Humic Materials



Leenheer et al. (1998)



Morel (1983)

Importance of Humic Materials

Global Carbon Reservoir

Take Part in Interfacial Phenomena

Undergo Coagulation and Aggregation

Involved in Photochemical Reactions

Contain Radicals

Known Reducing Agents

Methylate Metals

Form Chlorinated Species, THMs DBPs

Detoxify Metals

Limit Bioavailability of Metals

Alter Solubility

Influence Transport

Bind Metals & Organic Pollutants

Terminal Electron Acceptor for Bacteria

Humic material will aggregate & may “salt out” when it binds a cation




FIGURE 6. Structural model of a calcium inner-sphere complex

Leenheer, J.A. et al. (1998) *Environ. Sci. Technol.* 32, 2410

Importance of Humic Materials

Global Carbon Reservoir

Take Part in Interfacial Phenomena

Undergo Coagulation and Aggregation

Involved in Photochemical Reactions

Contain Radicals

Known Reducing Agents

Methylate Metals

Form Chlorinated Species, THMs DBPs

Detoxify Metals

Limit Bioavailability of Metals

Alter Solubility

Influence Transport

Bind Metals & Organic Pollutants

Terminal Electron Acceptor for Bacteria

Maturation and Fossilization are terms that refer to the formation of fossil fuels (coal, petroleum) from plant and animal material (biomolecules).

The overall process can be split into two or three major parts:

Marine → Diagenesis, Catagenesis, Metagenesis

Terrestrial → Humification, Coalification

May 1981, Volume 212, Number 4496

SCIENCE


Sludge Decomposition and Stabilization

Roy Hartenstein

of sludge decomposition and stabilization can be enhanced, to discuss the highly probable consequences of sludge stabilization in light of the basic information, and to suggest procedures for evaluating the sludge stabilization process. As a starting point, it is necessary to describe the fossilization pathway of the carbon cycle.

The Fossilization Pathway of the Carbon Cycle


Kerogens, coals, and petroleum oils are the earth's major fossil fuels; they



**Fossilization
Pathway of the
Carbon Cycle**
Hartenstein, 1981
Van Krevelen plot

Table 1. Analysis of organic Materials in Fossilization pathway (Percent dry wt.)									
	Carbo- hydrate	Protein	Fat	Mixed Diet	Sludge (act.)	Fulvic Acid	Humic Acid	Peat (old)	Coal (mid.)
C	44	58	75	53	32	47	59	59	85
H	6	7	12	7	4	4.4	5	6	5
N		11		2		2	3	2	1.5
O	49	23	12	36	37	46	34	31	8

from Hartenstein, 1981



Libes, 1992 "...diagenetic changes ...occur under anoxic conditions at temperatures less than 50 °C."



Pergamon

Applied Geochemistry, Vol. 11, pp. 711–720, 1996
Copyright © 1996 Elsevier Science Ltd
Printed in Great Britain. All rights reserved
0883–2927/96 \$15.00 + 0.00

Early diagenesis of organic matter in recent Black Sea sediments: characterization and source assessment

Abstract—The organic matter in 9 recent (not more than 250 years old) and 'organic-rich' sediments from the southern Black Sea shelf and upper slope have been characterized semi-quantitatively by Pyrolysis/Gas Chromatography/Mass Spectrometry (PY/GC/MS) and ¹³C Cross Polarization Magic Angle Spinning Nuclear Magnetic Resonance (CPMAS-NMR) spectrometry. The organic matter of 7 of the studied sediments was found to be ligno-carbohydrate with a proteinaceous component, one sediment appeared to contain oxidized coal dust and one contained thiophenes in association with pyrite. The ligno component is derived from grasses and soft wood lignin. Material entrapped in an anoxic environment contained the highest proportions of carbohydrate and protein. All the samples had suffered diagenesis as is generally shown by the attachment of carboxyl groups and the removal of methoxyl groups. The evidence suggests that diagenesis occurred whilst the particles traversed the oxic water column.

Sediment Diagenesis includes more than Organic Matter Transformations – Many redox processes occur






Table 12.2. *Stoichiometry of organic matter oxidation reactions*

Redfield ratios for x , y and z are 106, 16, 1.

Redox process	Reaction
Aerobic respiration	$(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4)_z + (x + 2y)\text{O}_2 \rightarrow$ $x\text{CO}_2 + (x + y)\text{H}_2\text{O} + y\text{HNO}_3 + z\text{H}_3\text{PO}_4$
Nitrate reduction	$5(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4)_z + 4x\text{NO}_3^- \rightarrow$ $x\text{CO}_2 + 3x\text{H}_2\text{O} + 4x\text{HCO}_3^- + 2x\text{N}_2 + 5y\text{NH}_3 + 5z\text{H}_3\text{PO}_4$
Manganese reduction	$(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4)_z + 2x\text{MnO}_2(\text{s}) + 3x\text{CO}_2 + x\text{H}_2\text{O} \rightarrow$ $2x\text{Mn}^{2+} + 4x\text{HCO}_3^- + y\text{NH}_3 + z\text{H}_3\text{PO}_4$
Iron reduction	$(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4)_z + 4x\text{Fe}(\text{OH})_3(\text{s}) + 7x\text{CO}_2 \rightarrow$ $4x\text{Fe}^{2+} + 8x\text{HCO}_3^- + 3x\text{H}_2\text{O} + y\text{NH}_3 + z\text{H}_3\text{PO}_4$
Sulfate reduction	$2(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4)_z + x\text{SO}_4^{2-} \rightarrow$ $x\text{H}_2\text{S} + 2x\text{HCO}_3^- + 2y\text{NH}_3 + 2z\text{H}_3\text{PO}_4$
Methane production	$(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4)_z \rightarrow$ $x\text{CH}_4 + x\text{CO}_2 + 2y\text{NH}_3 + 2z\text{H}_3\text{PO}_4$


Figure 12.1. A schematic representation of the porewater profiles that have been observed to show the sequential use of electron acceptors during organic matter degradation. Modified from Froelich *et al.* (1979).






Petroleum Maturation Process

Libes, 1992



Evolution of Fossil Fuels

Libes, 1992



Morel & Hering,
1993

Figure 6.13 A possible pathway for the formation of marine humic acids from a triglyceride. From Harvey et al., 1983.