

Math 192r, Problem Set #7 (Solutions)

1. (a) *How many different polygonal paths of length n are there that start at the point $(0, 0)$ and then take n steps of length 1, such that each step is either rightward, leftward, or upward, and such that no point gets visited more than once? Give an explicit formula.*

This is the same as the number of strings of length n consisting of the symbols R , L , and U (short for Right, Left, and Up, respectively) such that no R is followed by an L and no L is followed by an R . The associated 1-step transfer matrix

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$$

has characteristic polynomial $(t-1)(t^2-2t-1)$, so the answer is of the form $A+Br^n+Cs^n$ where $r = 1+\sqrt{2}$, $s = 1-\sqrt{2}$, and A, B, C are undetermined coefficients. Using the fact that the number of polygonal paths of the desired kind equals 1, 3, and 7 when n is 0, 1, and 2, respectively, we get $A = 0$, $B = 1+\sqrt{2}$, and $C = 1-\sqrt{2}$, so that the final answer is $\frac{1}{2}((1+\sqrt{2})^{n+1} + (1-\sqrt{2})^{n+1})$.

To do this in Maple, one might proceed as follows:

```
with(linalg):
m := matrix(3,3,[1,0,1,0,1,1,1,1,1]);
p := charpoly(m,t);
sols := solve(p=0,t);
r := sols[2];
s := sols[3];
ans:=solve({A+B*r+C*s=3,A+B*r^2+C*s^2=7,
           A+B*r^3+C*s^3=17},{A,B,C});
```

The result is a set whose three elements are equations giving the values of A , C , and B respectively. (Note that the Maple command I used doesn't return the values of the variables in the same order as I specified them! Does anyone know of a variant of my command that doesn't suffer from this defect?) By the way, the command `with(linalg)` only needs to be done once per session.

- (b) *If one chooses at random one of the paths of length n described in part (a) (so that each of the length- n paths has an equal chance of being chosen), what is the expected value of the y -coordinate of the last point on the path? Find a constant c so that this expected value is asymptotic to cn .*

An appropriate generating function is $\sum_{n \geq 1} (uM^n v)x^n$, where $u = (1, 1, y)$, $v = (1, 1, 1)^T$ (the transpose), and M is a modified version of the preceding transition matrix in which the 1's that correspond to Up-steps are replaced by y 's, so that when we multiply the matrix by itself, obtaining a matrix of polynomials in y , a term equal to y^k corresponds to a path that takes k Up-steps. (After we've expressed this generating function in closed form, we'll be able to differentiate it to get at the information we seek.) The entry y in the vector u occurs because it corresponds to taking a step in the Up direction. The generating function can be written as the sum of the nine entries of $u(\sum (Mx)^n)v = u(I - Mx)^{-1}v$, where the matrix Mx is

$$\begin{pmatrix} x & 0 & xy \\ 0 & x & xy \\ x & x & xy \end{pmatrix}.$$

We can use Maple for this:

```
with(linalg):
Mx := [[x,0,x*y],[0,x,x*y],[x,x,x*y]];
Id := [[1,0,0],[0,1,0],[0,0,1]];
u := [[x,x,x*y]];
v := [[1],[1],[1]];
inv := inverse(Id-Mx);
ans := simplify(multiply(u,inv,v)[1,1]);
```

(Note that for Maple, xy must be written as $x*y$; also note that Mx is just an indivisible symbol. Observe that the symbol I is reserved for the square root of minus 1. Finally, note that the output of `multiply(u,inv,v)` is a 1-by-1 matrix, not a number; hence the need to extract its 1,1 element with the matrix-entry-extraction operator `[1,1]`.) The answer `ans` turns out to be the

simple expression

$$\frac{2 + y + xy}{1 - x - xy - x^2y}.$$

If we differentiate this with respect to y and then set $y = 1$, we will obtain the generating function in which the coefficient of x^n is the sum of the heights of all the polygonal paths.

```
heights := simplify(subs(y=1,diff(ans,y)));
```

gives

$$\frac{x(1+x)^2}{(1-2x-x^2)^2}.$$

To find the asymptotic behavior of the coefficients of this generating function, use partial fractions over the field generated over the rationals by the square root of 2:

```
convert(heights, parfrac, x, sqrt(2));
```

Unfortunately, Maple gives us an answer in which the denominators of the four terms are of the form $x+a$ instead of $1+bx$, but this is only a minor annoyance. The term that controls the growth rate is the term whose denominator is quadratic and vanishes closest to $x = 0$. This is the term

$$\frac{1}{4} \frac{-1 + \sqrt{2}}{(x + 1 - \sqrt{2})^2} = \frac{1 + \sqrt{2}}{4} (1 - x(1 + \sqrt{2}))^{-2}.$$

Now we may apply the binomial theorem with exponent -2 : the coefficient of x^n in the preceding generating function equals

$$\frac{1 + \sqrt{2}}{4} \binom{-2}{n} (-(1 + \sqrt{2}))^n = \frac{1 + \sqrt{2}}{4} \binom{n+1}{n} (1 + \sqrt{2})^n,$$

which grows like $\frac{n}{4}(1 + \sqrt{2})^{n+1}$. The answer to part (a) grows like $\frac{1}{2}(1 + \sqrt{2})^{n+1}$, so, taking the ratio, we find that the expected height tends to the limit $n/2$. (There must be a nice way to see this!)

2. Look at the (infinite-dimensional) space of self-reciprocal Laurent polynomials in t , that is, Laurent polynomials in t that are unaffected by the substitution $t \rightarrow 1/t$. The space has two natural bases: $x_j = t^j + 1/t^j$ (with $j \geq 0$) and $y_k = (t + 1/t)^k$ (with $k \geq 0$).

- (a) Give an explicit formula for the entries of the (triangular) matrix that expresses the y_k in terms of the x_j .

I'll give the matrices in upper-triangular form (though I see that in the statement of the problem I used lower-triangular form, since I asserted in part (b) that each row has only finitely many non-zero entries). For $j = 0$, the j, k th entry in the matrix (the coefficient of x_j in the expansion of y_k) is $\frac{1}{2} \binom{k}{k/2}$ (which we interpret as 0 if $k/2$ is non-integer). For $j > 0$, the j, k th entry is simply $\binom{k}{(j+k)/2}$ (which we interpret as 0 if $k < j$ or if $(j+k)/2$ is not an integer). Here's the excerpt of the matrix with j, k running between 0 and 9:

$$\begin{pmatrix} 1 & 0 & 1 & 0 & 3 & 0 & 10 & 0 & 35 & 0 \\ 0 & 1 & 0 & 3 & 0 & 10 & 0 & 35 & 0 & 126 \\ 0 & 0 & 1 & 0 & 4 & 0 & 15 & 0 & 56 & 0 \\ 0 & 0 & 0 & 1 & 0 & 5 & 0 & 21 & 0 & 84 \\ 0 & 0 & 0 & 0 & 1 & 0 & 6 & 0 & 28 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 7 & 0 & 36 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 8 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 9 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

One way to make this array is as follows:

```
entry := proc(j,k)
  if (j=0 and k mod 2 = 0) then
    simplify((1/2)*binomial(k,k/2));
  elif (k>=j and j+k mod 2 = 0) then
    binomial(k,(j+k)/2);
  else 0; fi; end;
m := matrix(10,10,[seq(seq(entry(j,k),k=0..9),j=0..9)]);
```

- (b) Look at the inverse of this matrix (which expresses the x_j in terms of the y_k). Try to conjecture a formula for the entries, or if you

can't get that far, look for patterns and conjecture some interesting properties. (E.g., in each row of the infinite matrix, there are only finitely many non-zero values. What is their sum? What is the sum of their absolute values?) For full credit, you will need to conjecture (though not necessarily prove) a formula for all the entries.

In Maple (with the `linalg` library loaded), `inverse(m)` gives

$$\begin{pmatrix} 2 & 0 & -2 & 0 & 2 & 0 & -2 & 0 & 2 & 0 \\ 0 & 1 & 0 & -3 & 0 & 5 & 0 & -7 & 0 & 9 \\ 0 & 0 & 1 & 0 & -4 & 0 & 9 & 0 & -16 & 0 \\ 0 & 0 & 0 & 1 & 0 & -5 & 0 & 14 & 0 & -30 \\ 0 & 0 & 0 & 0 & 1 & 0 & -6 & 0 & 20 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & -7 & 0 & 27 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & -8 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & -9 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Column-sums are $2, 1, -1, -2, -1, 1, 2, 1, -1, -2, \dots$, which seems to be a periodic sequence. If we take absolute values before summing, we get $2, 1, 3, 4, 7, 11, 18, 29, 47, 76, \dots$, which looks like the Lucas sequence. We can certainly conjecture that the k, j entry has sign $(-1)^{(k-j)/2}$. (We index rows by k and columns by j now, since this is the inverse matrix.) One might also note that the successive non-zero diagonals (leaving aside the principal diagonal, which has a glitch in its first entry) appear to be given by polynomials of degree 1, 2, 3, etc.

The key to finding a formula for the entries lies in the observation that most of the entries in the j column are divisible by j , or nearly so. In fact, if you throw out the 0 row and 0 column, get rid of all the signs, and divide each entry by its column-index and then

multiply each entry by its row-index, you get the integer matrix

$$\begin{pmatrix} 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 2 & 0 & 3 & 0 & 4 & 0 \\ 0 & 0 & 1 & 0 & 3 & 0 & 5 & 0 & 10 \\ 0 & 0 & 0 & 1 & 0 & 4 & 0 & 10 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 5 & 0 & 15 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 6 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 7 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

which is just Pascal's triangle! So the coefficient of y_k in the expansion of x_j (relative to the basis y_0, y_1, \dots) is

$$(-1)^{(k-j)/2} \frac{j}{k} \binom{(j+k-2)/2}{(j-k)/2}$$

as long as $j \geq k \geq 1$ and j, k have the same parity; in the case $k = 0$, the coefficient is

$$(-1)^{(j-k)/2} 2$$

as long as $j \geq 0$ and j, k have the same parity; otherwise, the coefficient vanishes.