

Name \_\_\_\_\_

## GEOL3010L Earth Materials II Gravel Shape and Sedimentary Environment

### I. Introduction

Shape, size and composition are the fundamental properties of particles. Since shape is derived mostly by the processes of transport and deposition, studies of gravel assemblages can yield information on the environmental history of the sediments. Shape however is not easily measured or defined. Typically four elements of shape - form, sphericity, roundness and surface texture - are used to characterize particles.

**Form:** Form is a measure of the relative lengths of the three orthogonal axes of a clast. The longest axis (L), intermediate (I) and short axes (S) are measured with a vernier caliper. The oblate-prolate index (OP), determined by Dobkins and Folk (1970), ranges from  $-\infty$  to  $+\infty$ . The OP is a measure of the clast on a continuum from disc to blade to rod.

**Sphericity:** Sphericity measures the degree to which a particle approaches a spherical shape. It was defined by Wadell (1932) as the ratio between the diameter of a sphere with the same volume as the particle and the diameter of the circumscribed sphere.

**Roundness:** Roundness refers to the sharpness of the corners and edges of a grain. Roundness was defined by Wadell (1932) as the ratio of the average radius of curvature of the corners to the radius of the largest inscribed circle. Since it is quite time consuming to measure roundness, the common method of determining roundness is to visually compare grains of unknown roundness with standard images of grains of known roundness.

**Surface texture:** Surface texture is described by features found on the surface of the clast.

### II. Measurement of gravel clasts

For this exercise you should work in groups of two. One person should be responsible for estimating the roundness so that there is, hopefully, consistency in the roundness estimate. Select one of the clast trays. For most trays the clasts have been sorted on the basis of lithology.

**Identify the sample set on your lab (information requested on data sheet).**

1. Identify the rock type(s). What were the criteria you used to name the rock(s)? Be specific.

2. Determine the long (L), intermediate (I) and short (S) diameters of each of your pebbles using calipers. Enter the data on an Excel spreadsheet using the example below as a template. Do the necessary calculations.

No	D <sub>L</sub> (mm)	D <sub>I</sub> (mm)	D <sub>S</sub> (mm)	D <sub>S</sub> /D <sub>L</sub>	(D <sub>L</sub> - D <sub>I</sub> )/( D <sub>L</sub> - D <sub>S</sub> )	Form	D <sub>I</sub> /D <sub>L</sub>	D <sub>S</sub> /D <sub>I</sub>	Sphericity	Zingg Sphericity	Roundness
						<b>OP</b>			<b>Ψ<sub>p</sub></b>		
1	63.51	53.72	23.63	0.37	0.25	-6.84	0.85	0.44	0.55	Oblate	

3. Calculate **Form**: oblate-prolate index (OP) (Dobkins and Folk, 1970).

$$OP = \{10\{[(D_L - D_I) / (D_L - D_S)] - 0.5\} / (D_S/D_L)\}$$

4. Calculate **Sphericity**: maximum projection sphericity for each pebble, using Sneed and Folk's equation (Sneed and Folk, 1958).

$$\Psi_p = (D_S^2 / D_L D_I)^{1/3}$$

5. Classify the shape of each pebble using the Zingg diagram.

Plot each clast on the Zingg diagram

$$D_I/D_L \text{ vs } D_S/D_L$$

6. Estimate roundness using Krumbein's visual comparison chart.  
7. Calculate the mean and standard deviation of the form, sphericity and roundness for your clasts.


### III. Data Analysis

Make the following plots

- (1) A histogram showing the number of pebbles versus size (use the long axis measurement)
- (2) Clast size versus sphericity
- (3) Clast size versus roundness

The intervals for the histogram boxes will be determined after you have collected your data. For example, suppose your long axis dimensions vary from 10 to 50 mm, a size range of 40 mm. You want to have an adequate number of blocks so it might be convenient to divide the distribution into 8 blocks of 5mm each. In case you are not familiar with the construction of histograms an example is given here.

Size range	No. of Pebbles
10 – 15	2
15 – 20	5
20 – 25	8
25 – 30	10
30 – 35	14
35 – 40	11
40 – 45	7
45 – 50	3
Total	60



1. What is the size distribution of the clasts?
2. What is the most frequently appearing size?
3. Does size of clasts correlate with sphericity?
4. Does size of clasts correlate with roundness?

#### IV. Interpretation


When there is no correlation of grain size with sphericity or roundness (which indicates a process-controlled shape), you can use clast shape to determine the paleoenvironment. Specifically, using OP and  $\Psi_p$ , Dobkins and Folk (1970) determined the following for isotropic clasts within the size range of 16 - 126 mm:

- (i) Water as a transporting agent has a great effect on sphericity. In general fluvial gravel has  $\Psi_p \geq 0.67$  and beach gravels have  $\Psi_p \leq 0.65$ . Stratten (1974) and Gale (1990) found that mean  $\Psi_p$  for fluvial gravels is between 0.67 and 0.77 and for beach gravels between 0.53 and 0.64.
- (ii) Mean OP for fluvial gravel is near zero. Beach gravels have more negative OP (usually less than -2) and fluvial gravels are greater than -1.
- (iii) Using the mean Modified Wentworth Roundness scale, fluvial gravels range between 0.26 – 0.65. Low energy beach gravels (0.34 – 0.61) are less rounded than higher energy beach gravels (0.35-0.81).
- (iv) Glacial environments have been studied by Boulton (1978) and Dowdeswell et al. (1985). They found that plots of roundness versus sphericity can distinguish between the various transportation pathways.


For your clast set, determine the paleoenvironment it came from:

1. beach deposit
2. glacial environment
3. fluvial deposit
4. alluvial fan deposit

Compare your results with those of your classmates. Are your results the same? If not, how do they differ? If they differ what might be reason(s) for these differences?



Roundness from Krumbein (1941)



After Zingg (1935).