



ELSEVIER

Journal of Quantitative Spectroscopy &
Radiative Transfer ■ (■■■■) ■■■-■■■

Journal of
Quantitative
Spectroscopy &
Radiative
Transfer

www.elsevier.com/locate/jqsrt

Einstein A -coefficients and statistical weights for molecular absorption transitions in the *HITRAN* database

Marie Šimečková^{a,1}, David Jacquemart^{a,2}, Laurence S. Rothman^{a,*}, Robert
R. Gamache^b, Aaron Goldman^c

^aHarvard-Smithsonian Center for Astrophysics, Atomic and Molecular Physics Division, Cambridge, MA 02138-1516, USA

^bDepartment of Environmental, Earth & Atmospheric Sciences, University of Mass. Lowell, Lowell, MA 01854, USA

^cDepartment of Physics, University of Denver, Denver, CO 80208, USA

Received 28 August 2004; received in revised form 22 July 2005; accepted 22 July 2005

Abstract

This paper describes the calculation of the statistical weights and the Einstein A -coefficients for the 39 molecules and their associated isotopologues/isotopomers currently present in the line-by-line portion of the *HITRAN* database. Calculation of the Einstein A -coefficients was carried out using the *HITRAN* line intensities and the necessary statistical weights. The Einstein A -coefficient and the statistical weights of the upper and lower levels of the transition were added in the new format of the line parameters for the most recent edition of the *HITRAN* database.

© 2005 Published by Elsevier Ltd.

Keywords: Einstein A -coefficient; Statistical weight; Spectroscopic database; Atmospheric database; *HITRAN*

*Corresponding author. Tel.: +1 617 495 7474; fax: +1 617 496 7519.

E-mail address: lrothman@cfa.harvard.edu (L.S. Rothman).

¹New address: J. Heyrovský Institute of Physical Chemistry, Dolejškova 3, 18223 Praha 8, Czech Republic.

²New address: Laboratoire de Dynamiques, Interactions et Réactivités, Université Pierre et Marie Curie, 4 place Jussieu, Paris 75252, France.

1 1. Introduction

3 The *HITRAN* molecular spectroscopic database [1] consists of several components essential for
4 radiative transfer codes: high-resolution spectroscopic parameters of molecules in the gas phase,
5 absorption cross-sections for molecules with very dense spectral features, aerosol refractive
6 indices, ultraviolet line-by-line parameters and absorption cross-sections, and associated database
7 management software. This compilation can be accessed at [http://cfa-www.harvard.edu/
8 HITRAN](http://cfa-www.harvard.edu/HITRAN). The format of the line-by-line portion of the 2004 edition of *HITRAN* [1] has been
9 expanded. The statistical weights and the Einstein *A*-coefficients have been added for the
10 transitions of the thirty-nine molecules and their isotopologues/isotopomers³ (currently 93, see
11 Table 1 for the list). This paper discusses the motivation for adding these parameters and
12 describes their calculation in order to provide users with a clearer understanding.

13 Section 2 describes the motivation that led to the replacement of the weighted square of the
14 transition moment, \mathfrak{R} , that was present in the previous edition of *HITRAN* [2] by the Einstein *A*-
15 coefficient. In order to understand how the Einstein *A*-coefficients are calculated from the line
16 intensities given in *HITRAN*, the definition of the line intensity is presented in Section 3, and the
17 definition of the Einstein *A*-coefficient follows in Section 4. Section 5 gives the relation between
18 the line intensity and the Einstein *A*-coefficient. Section 6 clarifies the determination of the
19 statistical weights that are necessary to calculate the Einstein *A*-coefficients from the line
20 intensities. Furthermore, the determination of the so-called state-dependent nuclear spin
21 statistical weights is explained for each isotopologue/isotopomer present in the *HITRAN*
22 database. In Section 7, the relations between the Einstein *A*-coefficient, the oscillator strength, the
23 line intensity and the weighted square of the transition moment are recalled. Section 8 discusses
24 the non-additivity of the Einstein *A*-coefficients. Finally, Section 9 is devoted to inter-comparisons
25 between the Einstein *A*-values in this work and the Cologne database for molecular spectroscopy
26 [3].

29 2. Rationale for replacing the weighted square of the transition moment by the Einstein *A*-coefficient

31 The Einstein *A*-coefficient is desirable for applications in non-local thermodynamic equilibrium
32 (non-LTE) problems of the atmosphere, astrophysics, and fundamental physics. Reasons for
33 replacing the weighted square of the transition moment, \mathfrak{R} , by the Einstein *A*-coefficient in the
34 *HITRAN* database are discussed below.

35 As has been mentioned by Goldman et al. [4], several inconsistencies can occur with the use of
36 \mathfrak{R} . The first inconsistency is linked to the definition of the transition moment, which often differs
37 from one author to another. Indeed, the definition sometimes includes the Hönl–London factors,
38 sometimes not. For example, following Refs. [5–7], \mathfrak{R} includes the Hönl–London factors, as in the
39 definition for *HITRAN*. However in some studies [8,9], in order to reproduce the experimental *J*-

41 ³An isotopologue is a molecular variant that differs from the original molecule in the isotopic composition (number
42 of isotopic substitutions) only; for example, ¹³CH₄ and ¹²CH₃D are isotopologues of ¹²CH₄. An isotopomer (a
43 contraction of ‘isotopic isomer’), on the other hand, has the same number of each of the isotopic atoms but differing in
their orientation within the molecular structure (giving rise to different spectra), for example ¹⁶O¹⁸O¹⁶O is an
isotopomer of ¹⁶O¹⁶O¹⁸O and so is ¹⁴N¹⁵NO of ¹⁵N¹⁴NO.

1 dependence with the Herman Wallis factor, the Hönl–London factors were left out of \mathfrak{R} . This
 2 omission results in possible sources of error when data are incorporated into *HITRAN*.

3 Another inconsistency concerns the use of the statistical weights. In *HITRAN* [2], the parameter
 4 \mathfrak{R} is the weighted square of the transition moment as defined in Refs. [5,6]. However, because this
 5 definition is sometimes overlooked, or because the terminology is not precise enough (see for
 6 example Table 6 of Ref. [10]), the unweighted square of the transition moment has occasionally
 7 been put into previous editions of *HITRAN*. The nuclear spin statistical weights can also be
 8 included or left out in the definition of \mathfrak{R} . In order to be consistent with the definition of \mathfrak{R} with
 9 the Herman Wallis factor, these weights should not be included in the definition (see for example
 10 Jacquemart et al. [8]). Similarly, according to the convention adopted in Refs. [7,11], a
 11 supplemental weight equal to 2 can be introduced for the bands having *l*-type doubling. The fact
 12 that some authors neglect these supplemental weights, or use incorrect statistical weights (see for
 13 example Goldman et al. [4]), can lead to an erroneous value of \mathfrak{R} in *HITRAN*.

14 Another difficulty is related to the set of units used for \mathfrak{R} and to the expressions linking it to the
 15 *HITRAN* line intensity, which are different for electric-dipole, magnetic-dipole and electric-
 16 quadrupole transitions. The relation between the line intensity and \mathfrak{R} in Refs. [5–7] is given only
 17 for electric-dipole transitions. This equation is not rationalized [12], that is to say, the unit
 18 coherence of the equation is not respected since the vacuum permittivity, ϵ_0 , has been replaced by
 19 $1/4\pi$ in order to be in cgs electrostatic units. In *HITRAN*, the unit for \mathfrak{R} , which is valid for the
 20 electric-dipole transitions (the most prevalent cases in *HITRAN*), is Debye², but the standard unit
 21 is C² m². *HITRAN* also contains some magnetic-dipole transitions (O₂) and electric-quadrupole
 22 transitions (O₂ and N₂) for which the standard units are A² m⁴ and C² m⁴, respectively.

23 The choice of the Einstein *A*-coefficient is more standard and universal. There is only one
 24 definition and one unit for the Einstein *A*-coefficient (s⁻¹). Moreover, the Einstein *A*-coefficient
 25 does not depend on the type of a transition (i.e. electric-dipole, magnetic-dipole, electric-
 26 quadrupole...). The Einstein *A*-coefficient is also preferred to the Einstein *B*-coefficients because
 27 for *B* there are at least four possible cases according to the meaning given to the amount of
 28 radiation used in its definition [12,13].

29 It is important to point out the existence of a wide variety of expressions for the different
 30 quantities related to the interactions between molecules and light [13]. The whole formalism for
 31 the Einstein coefficients and the intensity can be found in Refs. [13,14]. However, in order to have
 32 the same notations and the same equations as in previous *HITRAN* papers [2,5,6,15], we
 33 summarize the most important steps that are necessary to obtain a relation between the Einstein
 34 *A*-coefficient and the line intensity.

37 3. Definition of the line intensity

38 According to the Lambert–Beer law, the variation $dI(\nu)$ of the intensity of a monochromatic
 39 radiance at wavenumber ν , going through a layer of a homogenous gas whose incremental depth is
 40 dL , is proportional to the depth of this layer and to the intensity $I(\nu)$ of this radiance [14]

$$43 \quad dI(\nu) = -k(\nu)I(\nu)dL, \quad (1)$$

where $k(\nu)$ (in cm⁻¹) is the spectral absorption coefficient per unit length. Note, the variables and

equations presented here adopt the notations used in *HITRAN* [2]; ν is the wavenumber⁴ and it is given in units of cm^{-1} . Care must be exercised when comparing the equations presented here with those found in other papers and texts where ν can sometimes be the frequency in Hz.

The integrated absorption coefficient S (in cm^{-2}) of an isolated line centered at ν_0 is defined as

$$S = \int_{-\infty}^{+\infty} k(\nu - \nu_0) d(\nu - \nu_0). \quad (2)$$

The spectral absorption coefficient can be expressed using S as a proportionality constant

$$k(\nu - \nu_0) = S\Phi(\nu - \nu_0), \quad (3)$$

since the integral of the normalized profile $\Phi(\nu - \nu_0)$ over the ~~all~~ spectral range is $\int_{-\infty}^{+\infty} \Phi(\nu - \nu_0) d(\nu - \nu_0) = 1$.

The absorption coefficient can be written as proportional either to the pressure P , or to the volumetric density ρ , or to the number N of absorbing molecules per unit volume

$$S = S_v^P P = S_v^\rho \rho = S_v^N N. \quad (4)$$

If P is in atm, ρ in g cm^{-3} , and N in molecule cm^{-3} , the units of S_v^P , S_v^ρ , and S_v^N are $\text{cm}^{-1} (\text{atm cm})^{-1}$, $\text{cm}^{-1} (\text{g cm}^{-2})^{-1}$, and $\text{cm}^{-1} (\text{molecule cm}^{-2})^{-1}$, respectively. Note that many other definitions and notations of the line intensity exist [16]. The notation of S_v^P , S_v^ρ and S_v^N is self-explanatory since the subscript ν means that we use wavenumbers (cm^{-1}), and the superscripts P , ρ , N mean that the line intensity is defined for a unit pressure, for a unit volumetric density, and for one molecule per unit volume, respectively.

It is important to note that in *HITRAN*, the given line intensity is S_v^N ($T = 296 \text{ K}$). Moreover, because S_v^P and S_v^N are the most common quantities used, it is useful to recall the relations between these two quantities

$$S_v^N(T) = \frac{P}{N} S_v^P(T). \quad (5)$$

For an ideal gas, we have

$$\frac{N}{P} = \frac{1}{k_B T} = \frac{1}{k_B T_S} \frac{T_S}{T}, \quad (6)$$

where $k_B = 1.38065 \times 10^{-16} \text{ erg K}^{-1}$ is the Boltzmann constant ($1 \text{ erg} = 10^{-7} \text{ J}$), T is the temperature of the gas, T_S is the standard temperature ($T_S = 273.15 \text{ K}$) and

$$\frac{1}{k_B T_S} = \frac{L(T_S)}{P_0}, \quad (7)$$

where $L(T_S)$ is the Loschmidt number ($L(T_S) = 2.68676 \times 10^{19} \text{ molecule cm}^{-3}$), and P_0 the pressure unit (equal to 1 atm). Finally, we obtain

$$S_v^N(T) = \frac{P_0}{L(T_S)} \frac{T}{T_S} S_v^P(T). \quad (8)$$

Eq. (8) provides the conversion between the two most common definitions of the line intensity.

⁴Throughout this paper we use the symbol ν for wavenumber (in units of cm^{-1}), rather than $\tilde{\nu}$ since there will be no confusion with frequency (units of s^{-1}).

The formula allowing the temperature conversion of the line intensity can be obtained by performing the ratio of the expressions of S_v^N (see Eq. (19) below and notations in Section 5) at two different temperatures, T and T'

$$S_v^N(T) = S_v^N(T') \frac{Q'_{\text{tot}}(T')}{Q_{\text{tot}}(T)} e^{-c_2 E_1((1/T)-(1/T'))} \left[\frac{1 - e^{-c_2 \nu/T}}{1 - e^{-c_2 \nu/T'}} \right], \quad (9)$$

where $Q_{\text{tot}}(T)$ is the total internal partition sum of the absorbing gas at the temperature T and E_1 is the energy of the lower state (cm^{-1}). c_2 is the second radiation constant, hc/k_B , where c is the speed of light and h is the Planck constant.

4. Definition of the Einstein coefficients

The Einstein A -coefficient for spontaneous emission is a first-order decay constant in units of s^{-1} [14]. If we consider that the number of atoms in the initial upper level m at time zero is $N_m(0)$, then, in the absence of an external radiation field, the number remaining after time t is

$$N_m(t) = N_m(0) e^{-\sum_n A_{mn} t}, \quad (10)$$

where the summation is performed over all the lower energy levels n .

Several definitions of the Einstein B -coefficients for induced absorption and emission exist, according to the meaning given to the amount of radiation of an external radiation field. In this work, the “amount of radiation”, u_ω , is taken as the energy density per unit frequency interval at the frequency of the line, ω , and it is expressed in $\text{J cm}^{-3} \text{ Hz}^{-1}$. Hence, the unit for B -coefficients is $\text{s}^{-1}/(\text{J cm}^{-3} \text{ Hz}^{-1})$, which is also written as $\text{cm}^3 (\text{J s}^2)^{-1}$. If we consider that the number of atoms in the initial level m is N_m , then the number of induced downward transitions to the lower level n per unit time is given by $N_m B_{mn} u_{\omega_{mn}}$. Similarly, the number of induced upward transitions to the upper level n' per unit time is $N_m B'_{mn} u_{\omega'_{mn}}$.

The rate at which N_m decreases with time as a result of these three radiative processes is described as

$$-\frac{dN_m}{dt} = N_m \sum_n A_{mn} + N_m \sum_n B_{mn} u_{\omega_{mn}} + N_m \sum_{n'} B'_{mn} u_{\omega'_{mn}} \quad (11)$$

(the replenishment of level m by transitions from other levels or its depletion or replenishment by collisional processes can be involved in further considerations).

In an approximation of a two level system (upper m and lower n levels are denoted as 2 and 1, respectively), we have the well-known equations linking the Einstein A - and B -coefficients

$$g_1 B_{12} = g_2 B_{21}, \quad (12)$$

$$A_{21} = 8\pi h \nu^3 B_{21}, \quad (13)$$

where A_{21} is in s^{-1} , and B_{12} and B_{21} are in $\text{cm}^3 (\text{J s}^2)^{-1}$, and g_1 and g_2 are the statistical weights of the levels 1 and 2, respectively. It is important to stress that Eq. (13) can be different if the units of the Einstein B -coefficients are different.

1 **5. Relation between the line intensity and the Einstein A -coefficient**

3 In order to obtain a relation between the line intensity and the Einstein A -coefficient, the first
 5 step is to express the variation of the intensity $dI(v - v_0)$ (see Eq. (1)) using the Einstein B -
 coefficients [14]

$$7 \quad dI(v - v_0) = -(N_1 B_{12} - N_2 B_{21}) \frac{h\nu_0}{c} I(v - v_0) \Phi(v - v_0) dl, \quad (14)$$

9 where N_2 and N_1 are the numbers of molecules per unit volume in the energy levels E_2 and E_1
 (expressed in cm^{-1}), respectively, such that $E_2 - E_1 = \nu_0$ is the wavenumber of the line.

11 The spectral absorption coefficient can be deduced from Eqs. (1) and (14):

$$13 \quad k(v - v_0) = (N_1 B_{12} - N_2 B_{21}) \frac{h\nu_0}{c} \Phi(v - v_0) \quad (15)$$

15 and the relation linking the line intensity to the Einstein B -coefficients is obtained from Eqs.
 (2)–(4), (15)

$$17 \quad S_v^N(T) = \frac{1}{N} \int_{-\infty}^{+\infty} k(v - v_0) d(v - v_0) = \frac{1}{N} (N_1 B_{12} - N_2 B_{21}) \frac{h\nu_0}{c}. \quad (16)$$

19 If N is the total number of molecules per unit volume at the temperature T , the population N_1 of
 the energy level E_1 is equal to

$$21 \quad N_1 = \frac{g_1 N}{Q_{\text{tot}}(T)} e^{-c_2 E_1 / T} \quad (17)$$

23 with a similar expression for N_2 . Combining Eqs. (16) and (17), we obtain

$$25 \quad S_v^N(T) = \frac{1}{Q_{\text{tot}}(T)} (e^{-c_2 E_1 / T} g_1 B_{12} - e^{-c_2 E_2 / T} g_2 B_{21}) \frac{h\nu_0}{c}. \quad (18)$$

27 Then using Eqs. (12) and (13), we have a relation between the line intensity and the Einstein A -
 29 coefficient

$$31 \quad S_v^N(T) = \frac{g_2}{Q_{\text{tot}}(T)} \frac{A_{21}}{8\pi c \nu_0^2} e^{-c_2 E_1 / T} (1 - e^{-c_2 \nu_0 / T}). \quad (19)$$

33 It is important to note that the line intensity given in these equations is defined for a pure gas.
 The line intensity in *HITRAN* is defined at the reference temperature $T_0 = 296$ K for an isotopic
 35 abundance, I_a . Calling this line intensity S_{HIT} , the conversion of the Einstein A_{21} coefficients from
 the line intensities S_{HIT} is performed according to

$$37 \quad A_{21} = \frac{8\pi c \nu_0^2 Q_{\text{tot}}(T_0) S_{\text{HIT}}}{e^{-c_2 E_1 / T_0} (1 - e^{-c_2 \nu_0 / T_0}) I_a g_2}. \quad (20)$$

39 Eq. (20) is valid for the electric-dipole, magnetic-dipole, and electric-quadrupole transitions that
 41 exist in *HITRAN*. It is equivalent to substituting Eq. (A.5) into Eq. (A.9) in Appendix A of Ref.
 [5]. However there is an error when similarly substituting Eq. (39) into Eq. (42) of Ref. [6] because
 43 the factor c had been omitted in Eq. (39). It is important to note that whereas the line intensities
 are additive, the Einstein A -coefficients are not (see Section 8).

In *HITRAN*, the conversion of the Einstein A_{21} coefficients from the line intensities S_{HIT} was performed using Eq. (20) with the recommended values of the physical constants c , h , and k_{B} (taken from Table XXIV of Mohr and Taylor [17]). All the other quantities, i.e. the lower state energy E_1 , the wavenumber of the transition ν_0 , the total internal partition function $Q_{\text{tot}}(T_0 = 296)$, the isotopic abundance I_a , the line intensity S_{HIT} , and the degeneracy factor g_2 of the upper level required in Eq. (20) are in the *HITRAN* database [1] either in the line-by-line portion (ν_0 , S_{HIT} , E_1 , and g_2), or in separate small files attached to the compilation ($Q_{\text{tot}}(T_0)$ and I_a). Table 1 summarizes the quantities $Q_{\text{tot}}(T_0)$ and I_a for each isotopologue present in *HITRAN*.

6. Calculation of the statistical weights

This section is devoted to the determination of the statistical weights that are required in the calculations of the Einstein A -coefficients from the line intensities (see Eq. (20)) and of the total internal partition functions of the molecules. In the *HITRAN* database, the partition functions come from the work of Refs. [4,18–20], and the statistical weights used in these studies agree with the definitions in Refs. [12,21–24]. The statistical weight g (also called the degeneracy factor) must be determined based on the quantum structure of the level:

$$g = g_e g_{\text{rot}}. \quad (21)$$

The factor g_e represents the total degeneracy due to all non-rotational factors, i.e. the product of the electronic statistical weight g_e and the vibrational statistical weight g_v , and the g_{rot} factor includes all the degeneracy factors related to the rotational state.

The statistical weight g of the level (g_2 of the upper level in Eq. (20)) is equal to g_{rot} for the transitions in *HITRAN*, which have the complete electronic, vibrational and rotational assignments (hyperfine assignments not considered, see Tables 3 and 4 of Ref. [1]). When the quantum numbers are not available in *HITRAN* (unassigned lines), zero values have been assigned for the statistical weights and likewise for the Einstein A -coefficients.

We will now consider several cases that appear in *HITRAN* requiring different formulae for the calculation of the statistical weights g_{rot} . In order to simplify the notations, the indices 1 and 2 for the lower and upper levels, respectively, are not reported in Eqs. (21)–(30).

6.1. Case where the hyperfine structure is not listed in *HITRAN*

For the transitions in *HITRAN* for which there is no F quantum number in the rotational field of the transition (see notations of local quanta identification in Ref. [1]), either because there is no hyperfine structure for the line or because it is not listed for the line, the statistical weight g_{rot} is equal to

$$g_{\text{rot}} = g_J = (2J + 1)g_s g_i, \quad (22)$$

where each state labeled by J has a $(2J + 1)$ degeneracy in the absence of an external electromagnetic field and g_s and g_i are the state-dependent and state-independent nuclear spin statistical weights, respectively.

The state-independent weight g_i is caused by the spins of the nuclei that are not interchanged in symmetry operations of a point group of a molecule [24]. It is given by the product over n nuclei of

1 Table 1
 2 Molecular entities in HITRAN with general quantities required to calculate the Einstein *A*-coefficient from the line
 3 intensity

5	Molecule	Isotopologue	State- independent weight, g_i	State- dependent weight, g_s	Total internal partition sum $Q_{\text{tot}}(T = 296 \text{ K})$	Fractional abundance I_a
7	1	H ₂ ¹⁶ O	1	1:3	1.7464E + 02	0.997317
9		H ₂ ¹⁸ O	1	1:3	1.7511E + 02	1.99983E – 03
		H ₂ ¹⁷ O	6	1:3	1.0479E + 03	3.71884E – 04
11		HD ¹⁶ O	6	1	8.5901E + 02	3.10693E – 04
		HD ¹⁸ O	6	1	8.7519E + 02	6.23003E – 07
		HD ¹⁷ O	36	1	5.2204E + 03	1.15853E – 07
13	2	¹² C ¹⁶ O ₂	1	1:0	2.8694E + 02	0.984204
15		¹³ C ¹⁶ O ₂	2	1:0	5.7841E + 02	1.10574E – 02
		¹⁶ O ¹² C ¹⁸ O	1	1	6.0948E + 02	3.94707E – 03
		¹⁶ O ¹² C ¹⁷ O	6	1	3.5527E + 03	7.33989E – 04
17		¹⁶ O ¹³ C ¹⁸ O	2	1	1.2291E + 03	4.43446E – 05
		¹⁶ O ¹³ C ¹⁷ O	12	1	7.1629E + 03	8.24623E – 06
19		¹⁸ O ¹² C ¹⁸ O	1	1:0	3.2421E + 02	3.95734E – 06
		¹⁷ O ¹² C ¹⁸ O	6	1	3.7764E + 03	1.47180E – 06
21	3	¹⁶ O ₃	1	1:0	3.4838E + 03	0.992901
		¹⁶ O ¹⁶ O ¹⁸ O	1	1	7.4657E + 03	3.98194E – 03
		¹⁶ O ¹⁸ O ¹⁶ O	1	1:0	3.6471E + 03	1.99097E – 03
23		¹⁶ O ¹⁶ O ¹⁷ O	6	1	4.3331E + 04	7.40475E – 04
		¹⁶ O ¹⁷ O ¹⁶ O	6	1:0	2.1405E + 04	3.70237E – 04
25	4	¹⁴ N ₂ ¹⁶ O	9	1	5.0018E + 03	0.990333
		¹⁴ N ¹⁵ N ¹⁶ O	6	1	3.3619E + 03	3.64093E – 03
27		¹⁵ N ¹⁴ N ¹⁶ O	6	1	3.4586E + 03	3.64093E – 03
		¹⁴ N ₂ ¹⁸ O	9	1	5.3147E + 03	1.98582E – 03
		¹⁴ N ₂ ¹⁷ O	54	1	3.0971E + 04	3.69280E – 04
29	5	¹² C ¹⁶ O	1	1	1.0712E + 02	0.986544
31		¹³ C ¹⁶ O	2	1	2.2408E + 02	1.10836E – 02
		¹² C ¹⁸ O	1	1	1.1247E + 02	1.97822E – 03
		¹² C ¹⁷ O	6	1	6.5934E + 02	3.67867E – 04
33		¹³ C ¹⁸ O	2	1	2.3582E + 02	2.22250E – 05
		¹³ C ¹⁷ O	12	1	1.3809E + 03	4.13292E – 06
35	6	¹² CH ₄	1	5:2:3	5.9045E + 02	0.988274
		¹³ CH ₄	2	5:2:3	1.1808E + 03	1.11031E – 02
		¹² CH ₃ D	3	8:4	4.7750E + 03	6.15751E – 04
37	7	¹⁶ O ¹⁶ O	1	1:0	2.1577E + 02	0.995262
		¹⁶ O ¹⁸ O	1	1	4.5230E + 02	3.99141E – 03
39		¹⁶ O ¹⁷ O	6	1	2.6406E + 03	7.42235E – 04
	8	¹⁴ N ¹⁶ O	3	1	1.1421E + 03	0.993974
41		¹⁵ N ¹⁶ O	2	1	7.8926E + 02	3.65431E – 03
		¹⁴ N ¹⁸ O	3	1	1.2045E + 03	1.99312E – 03
43	9	³² S ¹⁶ O ₂	1	1:0	6.3403E + 03	0.945678
		³⁴ S ¹⁶ O ₂	1	1:0	6.3689E + 03	4.19503E – 02

Table 1 (continued)

Molecule number	Isotopologue	State-independent weight, g_i	State-dependent weight, g_s	Total internal partition sum $Q_{\text{tot}}(T = 296 \text{ K})$	Fractional abundance I_a
10	$^{14}\text{N}^{16}\text{O}_2$	3	1:0	1.3578E + 04	0.991616
11	$^{14}\text{NH}_3$	3	4:2	1.7252E + 03	0.995872
	$^{15}\text{NH}_3$	2	4:2	1.1527E + 03	3.66129E – 03
12	$\text{H}^{14}\text{N}^{16}\text{O}_3$	6	1	2.1412E + 05	0.989110
13	^{16}OH	2	1	8.0362E + 01	0.997473
14	^{18}OH	2	1	8.0882E + 01	2.00014E – 03
15	^{16}OD	3	1	2.0931E + 02	1.55371E – 04
16	H^{19}F	4	1	4.1466E + 01	0.999844
17	H^{35}Cl	8	1	1.6066E + 02	0.757587
18	H^{37}Cl	8	1	1.6089E + 02	0.242257
19	H^{79}Br	8	1	2.0018E + 02	0.506781
20	H^{81}Br	8	1	2.0024E + 02	0.493063
21	H^{127}I	12	1	3.8900E + 02	0.999844
22	$^{35}\text{Cl}^{16}\text{O}$	4	1	3.2746E + 03	0.755908
23	$^{37}\text{Cl}^{16}\text{O}$	4	1	3.3323E + 03	0.241720
24	$^{16}\text{O}^{12}\text{C}^{32}\text{S}$	1	1	1.2210E + 03	0.937395
25	$^{16}\text{O}^{12}\text{C}^{34}\text{S}$	1	1	1.2535E + 03	4.15828E – 02
26	$^{16}\text{O}^{13}\text{C}^{32}\text{S}$	2	1	2.4842E + 03	1.05315E – 02
27	$^{16}\text{O}^{12}\text{C}^{33}\text{S}$	4	1	4.9501E + 03	7.39908E – 03
28	$^{18}\text{O}^{12}\text{C}^{32}\text{S}$	1	1	1.3137E + 03	1.87967E – 03
29	$\text{H}_2^{12}\text{C}^{16}\text{O}$	1	1:3	2.8467E + 03	0.986237
30	$\text{H}_2^{13}\text{C}^{16}\text{O}$	2	1:3	5.8376E + 03	1.10802E – 02
31	$\text{H}_2^{12}\text{C}^{18}\text{O}$	1	1:3	2.9864E + 03	1.97761E – 03
32	$\text{H}^{16}\text{O}^{35}\text{Cl}$	8	1	1.9274E + 04	0.755790
33	$\text{H}^{16}\text{O}^{37}\text{Cl}$	8	1	1.9616E + 04	0.241683
34	N_2	1	6:3	4.6598E + 02	0.992687
35	$\text{H}^{12}\text{C}^{14}\text{N}$	6	1	8.9529E + 02	0.985114
36	$\text{H}^{13}\text{C}^{14}\text{N}$	12	1	1.8403E + 03	1.10676E – 02
37	$\text{H}^{12}\text{C}^{15}\text{N}$	4	1	6.2141E + 02	3.62174E – 03
38	$^{12}\text{CH}_3^{35}\text{Cl}$	4	8:4	1.1583E + 05	0.748937
39	$^{12}\text{CH}_3^{37}\text{Cl}$	4	8:4	1.1767E + 05	0.239491
40	$\text{H}_2^{16}\text{O}_2$	1	1:3	9.8198E + 03	0.994952
41	$^{12}\text{C}_2\text{H}_2$	1	1:3	4.1403E + 02	0.977599
42	$^{12}\text{C}^{13}\text{CH}_2$	8	1	1.6562E + 03	2.19663E – 02
43	$^{12}\text{C}_2\text{H}_6$	1	24:20:16:8	7.0881E + 04	0.976990
44	PH_3	2	8:4	3.2486E + 03	0.999533
45	COF_2	1	1:3	7.0044E + 04	0.986544
46	SF_6	1	2:10:8:6	1.6233E + 06	0.950180
47	H_2^{32}S	1	1:3	5.0307E + 02	0.949884
48	H_2^{34}S	1	1:3	5.0435E + 02	4.21369E – 02
49	H_2^{33}S	4	1:3	2.0149E + 03	7.49766E – 03
50	HCOOH	4	1	3.9133E + 04	0.983898
51	HO_2	2	1	4.3004E + 03	0.995107

Table 1 (continued)

Molecule number	Isotopologue	State-independent weight, g_i	State-dependent weight, g_s	Total internal partition sum $Q_{\text{tot}}(T = 296 \text{ K})$	Fractional abundance I_a
34	O	1	See text	6.7212E + 00	0.997628
35	$^{35}\text{Cl}^{16}\text{O}^{14}\text{N}^{16}\text{O}_2$	12	1	4.7884E + 06	0.749570
	$^{37}\text{Cl}^{16}\text{O}^{14}\text{N}^{16}\text{O}_2$	12	1	4.9102E + 06	0.239694
36	NO^+	3	1	3.1168E + 02	0.993974
37	$\text{H}^{16}\text{O}^{79}\text{Br}$	8	1	2.8339E + 04	0.505579
	$\text{H}^{16}\text{O}^{81}\text{Br}$	8	1	2.8238E + 04	0.491894
38	$^{12}\text{C}_2\text{H}_4$	1	7:3	1.1041E + 04	0.977294
	$^{12}\text{C}^{13}\text{CH}_4$	2	10:6	4.5197E + 04	2.19595E – 02
39	CH_3OH	2	8:4	3.5314E + 04	0.985930

Note: All the values of the partition functions come from λ [18], except for the oxygen atom (see Section 6.3.34) and for CH_3OH [39].

2 times the nuclear spin I_n plus 1:

$$g_i = \prod_n (2I_n + 1). \quad (23)$$

The state-dependent weights g_s depend on the values of the total spin of the identical nuclei, which are interchanged in symmetry operations. Since, in this case, not all the nuclear spin states are always allowed to be associated with a specific rotational level, these weights vary from one transition to another and lead to the intensity alternation in the spectrum.

For example, the weights g_s of linear symmetrical molecules (point group $D_{\infty h}$) are different for the symmetric s (species Σ_g^+ and Σ_u^-) and antisymmetric a (species Σ_u^+ and Σ_g^-) rotational levels (see Fig. 99b of Ref. [24]). The following relation holds for those molecules with one pair of identical nuclei of spin I_x (see Eqs. (I.8) and (I.9) of Ref. [24, p. 17] for an arbitrary number of pairs of identical nuclei)

$$\begin{array}{l} \text{Fermi system — } s \text{ levels} \\ \text{Bose system — } a \text{ levels} \end{array} \quad g_s = \frac{1}{2}[(2I_x + 1)^2 - (2I_x + 1)], \quad (24a)$$

$$\begin{array}{l} \text{Fermi system — } a \text{ levels} \\ \text{Bose system — } s \text{ levels} \end{array} \quad g_s = \frac{1}{2}[(2I_x + 1)^2 + (2I_x + 1)]. \quad (24b)$$

This is the case of a molecule such as $^{12}\text{C}^{16}\text{O}_2$, which has two identical oxygen nuclei with spin zero. Substituting this value into the above equations yields a one-fold degeneracy for the even rotational J levels (symmetric s) and a zero-fold degeneracy for the odd rotational J levels (antisymmetric a) in the vibrationless electronic ground state (a totally symmetric state Σ_g^+).

For each isotopologue/isotopomer of each molecule in *HITRAN*, the values of g_i and g_s are summarized in Table 1. The description of the g_s calculation is also developed in Section 6.3.

6.2. Case where the hyperfine structure is listed in HITRAN

Hyperfine structure results from the intramolecular electromagnetic interactions of the nuclei with non-zero spins. For most of the transitions in *HITRAN*, the hyperfine structure is not resolved, that is to say, the hyperfine lines are not listed as separate transitions in the database. In these cases we have applied Eq. (22) to calculate the statistical weights g_{rot} . However, when the hyperfine structure is resolved (or partially resolved), Eq. (22) is no longer applicable.

Three cases altogether occur in *HITRAN* requiring different formulae for the calculation of the statistical weights g_{rot} , when the hyperfine structure is resolved. The first case, most common here, involves molecules such as $^{16}\text{O}^{12}\text{C}^{17}\text{O}$, $^{16}\text{O}^{17}\text{O}^{16}\text{O}$, $^{16}\text{O}^{16}\text{O}^{17}\text{O}$, $^{16}\text{O}^{17}\text{O}$, $^{14}\text{N}^{16}\text{O}$, $^{14}\text{N}^{16}\text{O}_2$, ^{16}OH , $^{35}\text{Cl}^{16}\text{O}$, $^{37}\text{Cl}^{16}\text{O}$, $^{12}\text{CH}_3^{35}\text{Cl}$, $^{12}\text{CH}_3^{37}\text{Cl}$, where there is only one nucleus that has non-zero spin (identical nuclei interchangeable in symmetry operations are not considered here). In the vector model, the \mathbf{I}_1 nuclear spin and \mathbf{J} molecular angular momenta are coupled together to form \mathbf{F}_1 , the total angular momentum of the molecule, which gives the splitting of each rotational J level into the hyperfine levels $F_1 = |J - I_1|, |J - I_1 + 1|, \dots, J + I_1$, each of them having $(2F_1 + 1)$ degeneracy in the absence of an external electromagnetic field. The statistical weight g_{rot} of the hyperfine level $F_1 (F_1 \equiv F)$ is equal to

$$g_{\text{rot}} = g_{F_1} = (2F_1 + 1)g_s \quad (25)$$

and the total number of Zeeman states is equal to the sum of the degeneracies of the hyperfine components

$$g_J = \sum_{F_1=|J-I_1|}^{J+I_1} g_{F_1} = g_s \sum_{F_1=|J-I_1|}^{J+I_1} (2F_1 + 1) = g_s(2J + 1)(2I_1 + 1) = (2J + 1)g_1g_s. \quad (26)$$

In this equation, the product in Eq. (23) reduces to $(2I_1 + 1)$.

The case where two nuclear spins are coupled together and to \mathbf{J} appears for a few molecules in *HITRAN* (some authors call this situation *hyperhyperfine* structure [21]). Examples in *HITRAN* include H^{79}Br , H^{81}Br , H^{35}Cl , H^{37}Cl , H^{127}I , and $\text{H}^{12}\text{C}^{14}\text{N}$. In the pure-rotation region, the hyperfine assignments for these molecules come from the JPL catalog [25]. The nuclear spin of Br, Cl, I, N, ... (denoted \mathbf{I}_1) is strongly coupled to \mathbf{J} to form \mathbf{F}_1 , and the spin of hydrogen (denoted \mathbf{I}_2) is weakly coupled to \mathbf{F}_1 to form \mathbf{F} . As a consequence, each J level is split into the levels $F_1 = |J - I_1|, |J - I_1 + 1|, \dots, J + I_1$, and further each F_1 level is split into the levels $F = |F_1 - I_2|, |F_1 - I_2 + 1|, \dots, F_1 + I_2$. In the JPL catalog (as in *HITRAN*), the quantum number associated with the hyperfine structure is just F_1 , not F (the hyperfine structure due to the hydrogen nucleus is not resolved at all). Note that there are no lines in the *HITRAN* database assigned with more than F_1 quantum number (we set $F_1 \equiv F$ for such cases).

The statistical weight g_{rot} of the hyperfine level F_1 in this case is equal to

$$g_{\text{rot}} = g_{F_1} = (2F_1 + 1)(2I_2 + 1)g_s \quad (27)$$

and the total number of Zeeman states is equal to the sum of the degeneracies over the states F_1

$$g_J = \sum_{F_1=|J-I_1|}^{J+I_1} g_{F_1} = g_s \sum_{F_1=|J-I_1|}^{J+I_1} (2F_1 + 1)(2I_2 + 1) = g_s(2J + 1)(2I_1 + 1)(2I_2 + 1) = (2J + 1)g_i g_s, \quad (28)$$

or to the sum of the degeneracies over the states F_1, F :

$$g_J = \sum_{F_1=|J-I_1|}^{J+I_1} \sum_{F=|F_1-I_2|}^{F_1+I_2} g_F = g_s \sum_{F_1=|J-I_1|}^{J+I_1} \sum_{F=|F_1-I_2|}^{F_1+I_2} (2F + 1) = g_s \sum_{F_1=|J-I_1|}^{J+I_1} (2F_1 + 1)(2I_2 + 1). \quad (29)$$

Fig. 1 represents a schematic example of the multiplets of HBr levels resulting from the coupling between \mathbf{J} and the nuclear spins \mathbf{I}_1 and \mathbf{I}_2 . In this figure, the statistical weights g_J , g_{F_1} , and g_F are indicated, allowing us to verify Eqs. (27)–(29).




Fig. 1. Schematic example of the multiplets of HBr levels.

The case of $\text{H}^{13}\text{C}^{14}\text{N}$ is a little more complicated, since we have three nuclear spins different from zero: $I(\text{H}) = I(^{13}\text{C}) = 1/2$, and $I(^{14}\text{N}) = 1$. In the *HITRAN* database (as in the JPL catalog), the F quantum number is in fact F_1 (F_1 is the result of the coupling of \mathbf{J} and $\mathbf{I}(^{14}\text{N})$). In this case, the statistical weight of the level F_1 is equal to

$$g_{\text{rot}} = g_{F_1} = (2F_1 + 1)(2I_2 + 1)(2I_3 + 1)g_s, \quad (30)$$

where I_2 is the nuclear spin of H, and I_3 of ^{13}C .

6.3. Determination of the statistical weights for each isotopologue/isotopomer

The statistical weights have been calculated for the upper and lower levels of each transition present in the *HITRAN* database [1]. The statistical weight, g , has been expressed as the product of g_e and g_{rot} (see Eq. (21)). For the calculation of g_{rot} we need to know the nuclear spin statistical weights g_i and g_s together with the quantum number J , where no hyperfine structure is resolved (Eq. (22)). Where the hyperfine structure due to the nucleus with the spin I_1 is resolved, the quantum number F_1 and the nuclear spin statistical weights $g_i/(2I_1 + 1)$ and g_s are needed (Eqs. (25), (27), (30)). The quantum numbers J and F_1 (denoted F in Table 4 of Ref. [1]) are given in the *HITRAN* line-by-line file of each molecule. The weights g_s and g_i are given in Table 1 for all the isotopologues present in *HITRAN*.

It follows from the selection rules that the nuclear spin statistical weights are the same for both the lower and upper levels of a transition. The state-dependent weights g_s are determined from the symmetries of the rotational levels (the electronic–vibration–rotation states) as described in the following subsections. In this text, the symmetries are the species of the point group of a molecule or the species of its rotational subgroup. In each subsection, we also indicate when hyperfine structure appears for an isotopologue, in order to determine which equation ((25), (27), (30)) has been used to calculate the statistical weights g_{rot} .

6.3.1. H_2O (molecule 1)

For the H_2^{16}O , H_2^{18}O , H_2^{17}O isotopologues (C_{2v}), the statistical weights g_s of the A and B rotational levels are 1 and 3, respectively. The symmetry of the rotational level is A if $K_a + K_c + v_3$ is even and B if $K_a + K_c + v_3$ is odd in the ground electronic state (the symmetry of the vibrational state is A if v_3 is even and B if v_3 is odd, see Figs. 17 and 144 of Ref. [24]). For the HD^{16}O , HD^{18}O , HD^{17}O isotopologues, the statistical weights g_s of all the rotational levels are 1.

No hyperfine structure is reported in the H_2O line list of the *HITRAN* database [1]. Consequently, Eq. (22) has been used to calculate the statistical weights g_{rot} for all the levels.

6.3.2. CO_2 (molecule 2)

For the $^{12}\text{C}^{16}\text{O}_2$, $^{13}\text{C}^{16}\text{O}_2$, $^{12}\text{C}^{18}\text{O}_2$ isotopologues ($D_{\infty h}$), the statistical weights g_s of the symmetric s and antisymmetric a rotational levels are 1 and 0, respectively (see Fig. 99 of Ref. [24], Fig. 17-6 of Ref. [26]). For the $^{16}\text{O}^{12}\text{C}^{18}\text{O}$, $^{16}\text{O}^{12}\text{C}^{17}\text{O}$, $^{16}\text{O}^{13}\text{C}^{18}\text{O}$, $^{16}\text{O}^{13}\text{C}^{17}\text{O}$, $^{17}\text{O}^{12}\text{C}^{18}\text{O}$ isotopologues, the statistical weights g_s of all the rotational levels are 1.

The statistical weights g_{rot} have been calculated using Eq. (25) for around 300 transitions of the $^{16}\text{O}^{12}\text{C}^{17}\text{O}$ isotopologue in the pure-rotation region, where the hyperfine structure is reported, and using Eq. (22) for all the other CO_2 lines in the *HITRAN* database.

6.3.3. O_3 (molecule 3)

For the $^{16}O^{16}O^{16}O$, $^{16}O^{17}O^{16}O$, $^{16}O^{18}O^{16}O$ isotopologues (C_{2v}), the statistical weights g_s of the A and B rotational levels are 1 and 0, respectively. The symmetry of the rotational level is A if $K_a + K_c + v_3$ is even and B if $K_a + K_c + v_3$ is odd in the ground electronic state (the symmetry of the vibrational state is A if v_3 is even and B if v_3 is odd, see Figs. 17 and 144 of Ref. [24]). For the $^{16}O^{16}O^{18}O$, $^{16}O^{16}O^{17}O$ isotopologues, the statistical weights g_s of all the rotational levels are 1.

In the current *HITRAN* edition [1], the hyperfine structure is reported for 78705 pure-rotational transitions of the $^{16}O^{17}O^{16}O$ and $^{16}O^{16}O^{17}O$ isotopomers in the spectral region between 0 and 156 cm^{-1} . The statistical weights g_{rot} have been calculated using Eq. (25) for the hyperfine lines and Eq. (22) for all the other ozone lines in the *HITRAN* database.

6.3.4. N_2O (molecule 4)

The statistical weights g_s of all the rotational levels are 1 for the nitrous oxide molecule.

No hyperfine structure is reported in the N_2O line list of the *HITRAN* database. Consequently, Eq. (22) has been used to calculate the statistical weights g_{rot} .

6.3.5. CO (molecule 5)

The statistical weights g_s of all the rotational levels are 1 for the carbon monoxide molecule.

No hyperfine structure is reported in the CO line list of the *HITRAN* database. Consequently, Eq. (22) has been used to calculate the statistical weights g_{rot} .

6.3.6. CH_4 (molecule 6)

For the $^{12}CH_4$ and $^{13}CH_4$ isotopologues (T_d), the statistical weights g_s of the A , E and F rotational levels are 5, 2 and 3, respectively [24,26]. The symmetry of the rotational level can be determined from the local quanta identification in *HITRAN*.

The symmetries of the rotational levels (and the statistical weights g_s) in the non-degenerate A and degenerate E vibrational states of the $^{12}CH_3D$ isotopologue (C_{3v}) in the electronic ground state are given in the Section 6.3.24. The $^{12}CH_3D$ isotopologue has three non-degenerate, $v_1(A_1)$, $v_2(A_1)$, $v_3(A_1)$, and three degenerate, $v_4(E)$, $v_5(E)$, $v_6(E)$, normal modes of vibration. The vibrational state has the A_1 symmetry if an arbitrary number of quanta of A_1 vibrations are excited ($\ell = 0$) and the E symmetry if one quantum of an E vibration is excited ($\ell = \pm 1$). If two quanta of an E vibration are excited, the vibrational state splits into the substates with the A_1 symmetry ($\ell = 0$) and the E symmetry ($\ell = \pm 2$). Similarly if one quantum of each of two different E vibrations are excited, the vibrational state splits into the substates with the $A_1 + A_2$ symmetry ($\ell = 0$) and the E symmetry ($\ell = \pm 2$). If $\ell = \pm 2$, each $K = |k| > 0$ rotational level is split into the so-called $+\ell$ and $-\ell$ levels (if these levels are assigned as $+\ell(k\ell < 0)$ and $-\ell(k\ell > 0)$, they have the same symmetries as if $\ell = \pm 1$, see Section 6.3.24).

No hyperfine structure is reported in the methane line list of the *HITRAN* database. Consequently, Eq. (22) has been used to calculate the statistical weights g_{rot} for all the methane lines.

6.3.7. O_2 (molecule 7)

For the $^{16}O^{16}O$ isotopologue ($D_{\infty h}$), the statistical weights g_s of the symmetric s and antisymmetric a rotational levels are 1 and 0, respectively (see Figs. 114b, 115b of Ref. [27]). For the $^{16}O^{18}O$ and $^{16}O^{17}O$ isotopologues, the statistical weights g_s of all the rotational levels are 1.

In the current *HITRAN* edition [1], the hyperfine structure is reported for the pure-rotational transitions of the $^{16}O^{17}O$ isotopologue in the spectral region between 0 and 251 cm^{-1} . The statistical weights g_{rot} have been calculated using Eq. (25) for the hyperfine lines and Eq. (22) for all the other oxygen lines in the *HITRAN* database.

6.3.8. NO (molecule 8)

The statistical weights g_s of all the rotational levels are 1 for the nitric oxide molecule. Hyperfine structure is reported in *HITRAN* for the transitions of the following bands of $^{14}N^{16}O$: ${}^2\Pi_{1/2}(v=1) \rightarrow {}^2\Pi_{1/2}(v=0)$, ${}^2\Pi_{3/2}(v=1) \rightarrow {}^2\Pi_{1/2}(v=0)$, ${}^2\Pi_{1/2}(v=1) \rightarrow {}^2\Pi_{3/2}(v=0)$, ${}^2\Pi_{3/2}(v=1) \rightarrow {}^2\Pi_{3/2}(v=0)$, ${}^2\Pi_{1/2}(v=2) \rightarrow {}^2\Pi_{1/2}(v=0)$, ${}^2\Pi_{3/2}(v=2) \rightarrow {}^2\Pi_{3/2}(v=0)$, ${}^2\Pi_{1/2}(v=2) \rightarrow {}^2\Pi_{1/2}(v=1)$, and ${}^2\Pi_{3/2}(v=2) \rightarrow {}^2\Pi_{3/2}(v=1)$. The statistical weights g_{rot} have been calculated using Eq. (25) for the hyperfine lines and Eq. (22) for all the other NO lines in the *HITRAN* database.

6.3.9. SO_2 (molecule 9)

For the $^{32}S^{16}O_2$ and $^{34}S^{16}O_2$ isotopologues (C_{2v}), the same rule holds as for the $^{16}O^{16}O^{16}O$ isotopologue (see Section 6.3.3).

No hyperfine structure is reported in the SO_2 line list of the *HITRAN* database. Consequently, Eq. (22) has been used to calculate the statistical weights g_{rot} for all the levels.

6.3.10. NO_2 (molecule 10)

For the $^{14}N^{16}O_2$ isotopologue (C_{2v}), the same rule holds as for the $^{16}O^{16}O^{16}O$ isotopologue (see Section 6.3.3).

Hyperfine structure is reported for all the NO_2 lines in *HITRAN* and the statistical weights g_{rot} have been calculated using Eq. (25).

6.3.11. NH_3 (molecule 11)

The ammonia molecule was for a long time classified among the pyramidal molecules (C_{3v}) with the normal modes of vibration, $v_1(A_1)$, $v_2(A_1)$, $v_3(E)$, $v_4(E)$, where the doublet levels are labeled s and a (symmetric and antisymmetric with respect to molecular inversion, respectively). In the (D_{3h}) labeling system, the normal modes of vibration are $v_1(A'_1)$, $v_2(A''_2)$, $v_3(E')$, $v_4(E')$, where the strongly anharmonic bending mode v_2 is defined as the inversion and the quantum number v_{inv} is introduced. There is the following correspondence between both the labeling systems: the $v_{\text{inv}} = 0$ (A'_1) corresponds to $v_2 = 0$ (s) ground state; $v_{\text{inv}} = 1$ (A''_2) corresponds to $v_2 = 0$ (a); $v_{\text{inv}} = 2$ (A'_1) corresponds to $v_2 = 1$ (s); $v_{\text{inv}} = 3$ (A''_2) corresponds to $v_2 = 1$ (a); etc. This means that the symmetry of the inversion state v_2 is A'_1 if v_{inv} is even (s state) and A''_2 if v_{inv} is odd (a state) where $v_{\text{inv}} = 0, 1$ if $v_2 = 0$; $v_{\text{inv}} = 2, 3$ if $v_2 = 1$, etc. [26].

If only non-degenerate vibrations are excited ($\ell = 0$), the symmetry of the vibrational state is A'_1 (s state) or A''_2 (a state). If one quantum of an E' vibration is excited ($\ell = \pm 1$) the symmetry of the vibrational state is E' (s state) or E'' (a state). If two quanta of an E' vibration are excited the

1 vibrational substates are s with the symmetries A'_1 ($\ell = 0$) and E' ($\ell = \pm 2$) or a with the
 2 symmetries A''_2 ($\ell = 0$) and E'' ($\ell = \pm 2$). Similarly if one quantum of each of two different E'
 3 vibrations are excited, the vibrational substates are s with the symmetries $A'_1 + A'_2$ ($\ell = 0$) and
 4 E' ($\ell = \pm 2$) or a with the symmetries $A''_2 + A''_1$ ($\ell = 0$) and E'' ($\ell = \pm 2$), see Sections 6.3.6 and
 5 6.3.24.

6 The rotational structures of the non-degenerate A and degenerate E vibrational states of the
 7 electronic ground state can be deduced from parts (a) and (b), respectively, of Fig. 118 of Ref. [24].
 8 The prime and double prime should be added over the symmetries of the rotational levels with the
 9 even and odd K , respectively, in the s states, and with the odd and even K , respectively, in the a
 10 states. Note that the A_1 and A_2 symmetries of the rotational levels in the A_2 vibrational states are
 11 interchanged (see also Table 15-1 and Fig. 15-5 of Ref. [26]).

12 The statistical weights g_s of the A'_2 , A''_2 and E' , E'' rotational levels are 4 and 2, respectively,
 13 while the A'_1 , A''_1 rotational levels are entirely missing [26,28].

14 No hyperfine structure is reported in the NH_3 line list of *HITRAN* and the statistical weights
 15 g_{rot} have been calculated using Eq. (22).

17 6.3.12. HNO_3 (molecule 12)

18 The statistical weights g_s of all the rotational levels are 1 for the nitric acid molecule.

19 No hyperfine structure is reported in the HNO_3 line list of *HITRAN* and the statistical weights
 20 g_{rot} have been calculated using Eq. (22).

22 6.3.13. OH (molecule 13)

23 The statistical weights g_s of all the rotational levels are 1 for the hydroxyl radical, OH.

24 The statistical weights g_{rot} of the hyperfine lines in the pure-rotation region have been
 25 calculated using Eq. (25); otherwise Eq. (22) has been used where no hyperfine structure is
 26 reported.

28 6.3.14. HF (molecule 14)

29 The statistical weights g_s of all the rotational levels are 1 for the hydrogen fluoride molecule.

30 Hyperfine structure is not reported in the HF line list of *HITRAN*. Eq. (22) has been used to
 31 calculate the statistical weights g_{rot} .

33 6.3.15. HCl (molecule 15)

34 The statistical weights g_s of all the rotational levels are 1 for the hydrogen chloride molecule.

35 For the hyperfine lines of H^{35}Cl and H^{37}Cl in the pure-rotation region, Eq. (27) has been used
 36 to calculate the statistical weights g_{rot} . Let us recall that for these levels, the hyperfine quantum
 37 number F given in *HITRAN* is F_1 (the nuclear spin of Cl (denoted I_1) is strongly coupled to \mathbf{J} to
 38 form \mathbf{F}_1 , see Section 6.2). For all the other lines, where no hyperfine structure is reported, Eq. (22)
 39 has been used.

41 6.3.16. HBr (molecule 16)

42 The statistical weights g_s of all the rotational levels are 1 for the hydrogen bromide molecule.

43 For the hyperfine lines of H^{79}Br and H^{81}Br of the (0–0) and (1–0) bands, Eq. (27) has been used
 to calculate the statistical weights g_{rot} . For these levels, the hyperfine quantum number F given in

HITRAN is F_1 (the nuclear spin of Br (denoted I_1) is strongly coupled to J to form F_1 , see Section 6.2). For all the other lines, where no hyperfine structure is reported, Eq. (22) has been used.

6.3.17. HI (molecule 17)

The statistical weights g_s of all the rotational levels are 1 for the hydrogen iodide molecule.

For the hyperfine lines of $H^{127}I$ of the (0–0) and (1–0) bands, Eq. (27) has been used to calculate the statistical weights g_{rot} . For these levels, the hyperfine quantum number F given in *HITRAN* is F_1 (the nuclear spin of I (denoted I_1) is strongly coupled to J to form F_1 , see Section 6.2). For all the other lines, where no hyperfine structure is reported, Eq. (22) has been used.

6.3.18. ClO (molecule 18)

The statistical weights g_s of all the rotational levels are 1 for the chlorine monoxide molecule.

Hyperfine structure is reported in *HITRAN* for the transitions of the ${}^2\Pi_{1/2}(v=0) - {}^2\Pi_{1/2}(v=0)$, ${}^2\Pi_{3/2}(v=0) - {}^2\Pi_{3/2}(v=0)$ bands of ${}^{35}\text{Cl}^{16}\text{O}$ and ${}^{37}\text{Cl}^{16}\text{O}$. The statistical weights g_{rot} of the hyperfine levels have been calculated from Eq. (25), otherwise Eq. (22) has been used where no hyperfine structure is reported.

6.3.19. OCS (molecule 19)

The statistical weights g_s of all the rotational levels are 1 for the carbonyl sulfide molecule.

Hyperfine structure is not reported in the OCS line list of *HITRAN*. Eq. (22) has been used to calculate the statistical weights g_{rot} .

6.3.20. H₂CO (molecule 20)

For the H₂CO molecule (C_{2v}), the statistical weights g_s of the A and B rotational levels are 1 and 3, respectively. The symmetry of the rotational level is A if $K_a + v_3 + v_4 + v_5$ is even and B if $K_a + v_3 + v_4 + v_5$ is odd in the ground electronic state (the symmetry of the vibrational state is A if $v_3 + v_4 + v_5$ is even and B if $v_3 + v_4 + v_5$ is odd, see Figs. 17 and 143 of Ref. [24]).

Hyperfine structure is not reported in the H₂CO line list of *HITRAN*; thus Eq. (22) has been used to calculate the statistical weights g_{rot} .

6.3.21. HOCl (molecule 21)

The statistical weights g_s of all the rotational levels are 1 for the hypochlorous acid molecule.

No hyperfine structure is reported in the HOCl line list of the *HITRAN* database. Consequently, Eq. (22) has been used to calculate the statistical weights g_{rot} .

6.3.22. N₂ (molecule 22)

For the nitrogen molecule ($D_{\infty h}$) in the ground electronic state, the statistical weights g_s of the symmetric s (even J) and antisymmetric a (odd J) rotational levels are 6 and 3, respectively (see Figs. 64b, 114b of Ref. [27]).

No hyperfine structure is reported in the N₂ line list of the *HITRAN* database. Consequently, Eq. (22) has been used to calculate the statistical weights g_{rot} .

6.3.23. HCN (molecule 23)

The statistical weights g_s of all the rotational levels are 1 for the hydrogen cyanide molecule.

1 The statistical weights g_{rot} of the hyperfine components of the $R(0)$ lines in the pure-rotational
 2 bands (000–000) of $\text{H}^{12}\text{C}^{14}\text{N}$ and $\text{H}^{13}\text{C}^{14}\text{N}$ were calculated from Eqs. (27) and (30), respectively.
 3 Let us recall that for these levels, the hyperfine quantum number F given in *HITRAN* is F_1 (the
 4 nuclear spin of ^{14}N (denoted \mathbf{I}_1) is strongly coupled to \mathbf{J} to form \mathbf{F}_1 , see Section 6.2). For all the
 5 other lines, where no hyperfine structure is reported, Eq. (22) has been used.

7 6.3.24. CH_3Cl (molecule 24)

9 The symmetries A and E of the rotational levels of a C_{3v} molecule in a totally symmetric
 10 electronic state are determinable from the values of the quantum number $G = \ell - k$ introduced by
 11 Hougen [29] ($\ell = \sum_i \ell_i$ runs over all degenerate normal modes, where $\ell_i = v_i, v_i - 2, \dots, -v_i$ is the
 12 vibrational angular momentum quantum number associated with the i th mode, and
 13 $k = 0, \pm 1, \pm 2, \dots, \pm J$). The rotational level has the $A_1 + A_2$ symmetry if $G = 0, \pm 3, \pm 6, \pm 9, \dots$
 14 (an exception occurs if $k = 0$ and $\ell = 0$ see below) and the E symmetry if $G = \pm 1, \pm 2, \pm 4, \pm 5, \dots$.
 15 The statistical weights g_s of the A_1, A_2 and E rotational levels are 4 (the weight is 8 for the $A_1 + A_2$
 16 level) [26,28]. The symmetry of the rotational level has been added in the local quanta field of the
 17 current *HITRAN* edition, but only for the recent work between 6 and $8\ \mu\text{m}$ [30].

18 The CH_3Cl molecule has three non-degenerate, $v_1(A_1), v_2(A_1), v_3(A_1)$, and three degenerate,
 19 $v_4(E), v_5(E), v_6(E)$, normal modes of vibration. If an arbitrary number of quanta of A_1 vibrations
 20 are excited ($\ell = 0$) the symmetry of the vibrational state is A_1 and the symmetries of the rotational
 21 levels are $A_1 + A_2$ if $K = 3, 6, 9, \dots$ and E if $K = 1, 2, 4, 5, \dots$; the $K = 0$ level has the symmetry A_1
 22 if J is even and A_2 if J is odd (see Figs. 118a of Ref. [24], 9.1A of Ref. [28]).

23 If one quantum of an E vibration is excited ($\ell = \pm 1$) the symmetry of the vibrational state is E
 24 and each $K = |k| > 0$ rotational level is split into the so-called $+\ell$ level ($k\ell > 0$ and
 25 $G = \pm(|\ell| - |k|)$) and $-\ell$ level ($k\ell < 0$ and $G = \pm(|\ell| + |k|)$). The $+\ell$ and $-\ell$ levels have the
 26 symmetries $A_1 + A_2$ and E , respectively, if $K = 1, 4, \dots$; E and $A_1 + A_2$, respectively, if $K =$
 27 $2, 5, \dots$; E and E , respectively, if $K = 3, 6, \dots$; and the symmetry is E if $K = 0$ (see Figs. 118b of
 28 Ref. [24], 9.1B of Ref. [28]).

29 If three quanta of an E vibration are excited, the vibrational state splits into the substates with
 30 the $A_1 + A_2$ symmetry ($\ell = \pm 3$) and the E symmetry ($\ell = \pm 1$). The symmetries of the rotational
 31 levels in the A_1 and E vibrational states are given above. (The rotational structure of the A_2
 32 vibrational state is similar to that of the A_1 vibrational state except that the A_1 and A_2 symmetries
 33 of the rotational levels are interchanged.)

34 Hyperfine structure is reported in *HITRAN* for the transitions of the pure-rotational bands of
 35 $^{12}\text{CH}_3^{35}\text{Cl}$ and $^{12}\text{CH}_3^{37}\text{Cl}$. The statistical weights g_{rot} have been calculated using Eq. (25) for the
 36 hyperfine lines and Eq. (22) for all the other lines.

37 6.3.25. H_2O_2 (molecule 25)

38 For the hydrogen peroxide molecule (C_2), the statistical weights g_s of the A and B rotational
 39 levels are 1 and 3, respectively. The symmetry of the rotational level is A if K_c is even and B if K_c is
 40 odd in the ground and $v_3 = 1$ vibrational states (A symmetry), while the reverse is true for the
 41 $v_6 = 1$ vibrational state (B symmetry) of the electronic ground state. These rules hold
 42 independently of the values of the torsional quantum numbers n and τ (see note of Table 3 of
 43 Ref. [1]).

1 No hyperfine structure is reported in the H_2O_2 line list of the *HITRAN* database. Consequently,
 2 Eq. (22) has been used to calculate the statistical weights g_{rot} .

3 6.3.26. C_2H_2 (molecule 26)

4 For the $^{12}\text{C}_2\text{H}_2$ isotopologue ($\mathbf{D}_{\infty\text{h}}$), the statistical weights g_s of the symmetric s and
 5 antisymmetric a rotational levels are 1 and 3, respectively. The symmetries s/a of the rotational
 6 levels of a given vibrational state ($\Sigma_{g(u)}^+$, $\Sigma_{g(u)}^-$, $\Pi_{g(u)}$, $\Delta_{g(u)}$) of the electronic ground state can be
 7 deduced from Fig. 99 of Ref. [24] (Fig. 17-6 of Ref. [26]).

8 The rotational levels in *HITRAN* are assigned as e and f , respectively: the e levels are those with
 9 the parity $+(-1)^J$ and the f levels are those with the parity $-(-1)^J$ [26,31]. The symmetry of the
 10 vibrational state can be determined from the local quanta identification in *HITRAN* (note that the
 11 characters u (ungerade) and g (gerade) have been added in the new vibrational format, see Table 3
 12 of Ref. [1]).

13 For the $^{12}\text{C}^{13}\text{CH}_2$ isotopologue, the statistical weights g_s of all the rotational levels are 1.

14 No hyperfine structure is reported in the C_2H_2 line list of the *HITRAN* database and Eq. (22)
 15 has been used to calculate the statistical weights g_{rot} .

16 6.3.27. C_2H_6 (molecule 27)

17 The symmetries of the rotational levels of the ethane molecule (staggered model, $\mathbf{D}_{3\text{d}}$ point
 18 group) in the ground electronic state can be deduced from Fig. 118a of Ref. [24] for the A_1 and A_2
 19 vibrational states (note that the A_1 and A_2 symmetries of the rotational levels in the A_2 vibrational
 20 states are interchanged) and from Fig. 118b of Ref. [24] for the E vibrational states. All the
 21 rotational levels are g and u levels in the g and u vibrational states, respectively. The statistical
 22 weights g_s of the A_1 , A_2 and E rotational levels are 8, 16 and 20, respectively [26,32].

23 The rotational levels of the vibrational ground state (A_{1g} symmetry) have the symmetries
 24 $A_{1g} + A_{2g}$ if $K = 3, 6, 9, \dots$ (the weight factor is 24); E_g if $K = 1, 2, 4, 5, \dots$ (the weight factor is
 25 20); A_{1g} if $K = 0$ and J is even (the weight factor is 8); A_{2g} if $K = 0$ and J is odd (the weight factor
 26 is 16). All these levels change from g to u if an odd number of torsional quanta is excited ($v_4 = 1$
 27 vibrational state, A_{1u} symmetry).

28 In the $v_7 = 1$ vibrational state (E_u symmetry), the so-called $+\ell$ and $-\ell$ levels ($\ell = \pm 1$, see
 29 Section 6.3.24) have the symmetries $A_{1u} + A_{2u}$ and E_u , respectively, if $K = 1, 4, \dots$; E_u and
 30 $A_{1u} + A_{2u}$, respectively, if $K = 2, 5, \dots$; E_u and E_u , respectively, if $K = 3, 6, \dots$; and the symmetry
 31 is E_u if $K = 0$ (the weight factor is 24 for $A_{1u} + A_{2u}$ and 20 for E_u levels). All these levels change
 32 from u to g if an odd number of torsional quanta is excited ($v_7 = 1$, $v_4 = 1$ vibrational state, E_g
 33 symmetry).

34 If the internal rotation tunneling leads to observable splitting in the spectrum, the rotational
 35 levels (electronic–vibration–rotation–torsion states) have the symmetry species s of the \mathbf{G}_{36}^+
 36 permutation-inversion group: $A_{1g}(8) \rightarrow A_{1s}(6) + E_{3s}(2)$, $A_{2g}(16) \rightarrow A_{2s}(10) + E_{4s}(6)$, $E_g(20) =$
 37 $E_{1s}(4) + G_s(16)$ and $A_{1u}(8) \rightarrow A_{3s}(6) + E_{3s}(2)$, $A_{2u}(16) \rightarrow A_{4s}(10) + E_{4s}(6)$,
 38 $E_u(20) = E_{2s}(4) + G_s(16)$, the statistical weights g_s are given in parentheses [33]
 39 ($A_{1g} + A_{2g} \rightarrow A_s(16) + E_s(8)$ in the *HITRAN* file where the splitting is only partially resolvable).

40 No hyperfine structure is reported in the C_2H_6 line list of the *HITRAN* database and Eq. (22)
 41 has been used to calculate the statistical weights g_{rot} .

6.3.28. PH_3 (molecule 28)

The symmetries of the vibrational states of the phosphine molecule, which has four normal modes of vibration, $v_1(A_1)$, $v_2(A_1)$, $v_3(E)$, $v_4(E)$, and the symmetries of the rotational levels (and the statistical weights g_s) in the non-degenerate A and degenerate E vibrational states of the electronic ground state are determinable under the rules given for C_{3v} molecules in the Sections 6.3.6 and 6.3.24.

No hyperfine structure is reported in the PH_3 line list of the *HITRAN* database. Consequently, Eq. (22) has been used to calculate the statistical weights g_{rot} .

6.3.29. COF_2 (molecule 29)

For the carbonyl fluoride molecule (C_{2v}), the same rule holds as for the formaldehyde molecule (see Section 6.3.20).

No hyperfine structure is reported in the COF_2 line list of the *HITRAN* database. Consequently, Eq. (22) has been used to calculate the statistical weights g_{rot} .

6.3.30. SF_6 (molecule 30)

The statistical weights g_s of the A_1 , A_2 , E , F_1 and F_2 rotational levels of the sulfur hexafluoride molecule (O_h) are 2, 10, 8, 6 and 6, respectively [26,32,34]. This symmetry appears in the rotational assignment of *HITRAN* (designated by assignment C in *HITRAN*, see Table 4 of Ref. [1]).

No hyperfine structure is reported in the SF_6 line list of the *HITRAN* database and Eq. (22) has been used to calculate the statistical weights g_{rot} for all the levels.

6.3.31. H_2S (molecule 31)

For the $H_2^{32}S$, $H_2^{33}S$ and $H_2^{34}S$ isotopologues (C_{2v}), the same rule holds as for the $H_2^{16}O$ isotopologue (see Section 6.3.1).

No hyperfine structure is reported in the H_2S line list of the *HITRAN* database. Consequently, Eq. (22) has been used for all the levels to calculate the statistical weights g_{rot} .

6.3.32. $HCOOH$ (molecule 32)

The statistical weights g_s of all the rotational levels are 1 for the formic acid molecule.

No hyperfine structure is reported in the $HCOOH$ line list of the *HITRAN* database and Eq. (22) has been used to calculate the statistical weights g_{rot} .

6.3.33. HO_2 (molecule 33)

The statistical weights g_s of all the rotational levels are 1 for the hydroperoxy radical, HO_2 .

Hyperfine structure is reported in *HITRAN* for the transitions of the (000–000) band of HO_2 . The statistical weights g_{rot} have been calculated using Eq. (25) for the hyperfine lines and Eq. (22) for all the other lines.

6.3.34. O (molecule 34)

The oxygen atom is unique in the database, having no vibration-rotation states. Only two electronic transitions are present in *HITRAN*: the first one is at 68.716 cm^{-1} between the upper level $2s^22p^4\ ^3P_0$ and the lower level $2s^22p^4\ ^3P_1$, the second is at 158.303 cm^{-1} between the upper level $2s^22p^4\ ^3P_1$ and the lower level $2s^22p^4\ ^3P_2$. The statistical weight, which appears in Eq. (20), is

equal to 5 for the level $2s^22p^4\ ^3P_2$, 3 for the level $2s^22p^4\ ^3P_1$, and 1 for the level $2s^22p^4\ ^3P_0$. The electronic partition sum used to obtain the Einstein A -coefficients from the *HITRAN* line intensities is equal to 6.72123. It has been calculated as usual

$$Q = \sum_i g_i e^{-c_2 E_i/T}, \quad (31)$$

where g_i and E_i are the statistical weight and the energy of the level i , respectively. The summation is performed over all the possible states i . In our case, the summation has been performed over the three P states.

No hyperfine structure is reported in the oxygen-atom line list of the *HITRAN* database. Consequently, Eq. (22) has been used for all the levels to calculate the statistical weights g_{rot} .

The calculated values of the Einstein A -coefficients in this work have been compared with the values in the literature. For the 68.716 cm^{-1} transition, we obtain $1.74 \times 10^{-5}\text{ s}^{-1}$, as compared to $1.78 \times 10^{-5}\text{ s}^{-1}$ from Galavís et al. [35]; for the 158.303 cm^{-1} transition, we obtain $8.80 \times 10^{-5}\text{ s}^{-1}$, as compared to $8.86 \times 10^{-5}\text{ s}^{-1}$ from Ref. [35] and to $8.46 \times 10^{-5}\text{ s}^{-1}$ from Vastel et al. [36].

6.3.35. ClONO_2 (molecule 35)

The statistical weights g_s of all the rotational levels are 1 for the chlorine nitrate molecule.

No hyperfine structure is reported in the ClONO_2 line list of the *HITRAN* database and Eq. (22) has been used for all the levels to calculate the statistical weights g_{rot} .

6.3.36. NO^+ (molecule 36)

The statistical weights g_s of all the rotational levels are 1 for the NO^+ cation.

No hyperfine structure is reported in the NO^+ line list of the *HITRAN* database. Consequently, Eq. (22) has been used for all the levels to calculate the statistical weights g_{rot} .

6.3.37. HOBr (molecule 37)

The statistical weights g_s of all the rotational levels are 1 for the hypobromous acid molecule.

No hyperfine structure is reported in the HOBr line list of the *HITRAN* database and Eq. (22) has been used for all the levels to calculate the statistical weights g_{rot} .

6.3.38. C_2H_4 (molecule 38)

For the $^{12}\text{C}_2\text{H}_4$ isotopologue (\mathbf{D}_{2h}), the statistical weights g_s of the A , B_1 , B_2 and B_3 rotational levels are 7, 3, 3, and 3, respectively. The symmetries of the rotational levels in the A , B_1 , B_2 and B_3 vibrational states of the electronic ground state can be deduced from Figs. 145a–d of Ref. [24], respectively. (Note, that Herzberg's convention of the symmetry labeling of the normal modes of vibration is adopted here, see Fig. 44 of Ref. [24] and Table 6 of Ref. [37]).

In the ground vibrational state (A symmetry), the symmetry of the rotational level is A if both K_a and K_c are even, and B if K_a or K_c is odd (see Figs. 17 and 145a of Ref. [24]); in the $v_{10} = 1$ vibrational state (B_2 symmetry), the symmetry of the rotational level is A if both K_a and K_c are odd and B if K_a or K_c is even (see Figs. 17 and 145c of Ref. [24]) [38].

For the $\text{H}_2^{12}\text{C}^{13}\text{CH}_2$ isotopologue (\mathbf{C}_{2v}), the statistical weights g_s of the A and B rotational levels are 10 and 6, respectively. In the ground vibrational state (A symmetry), the symmetry of the rotational level is A if K_a is even and B if K_a is odd (see Figs. 17 and 143a of Ref. [24]).

1 No hyperfine structure is reported in the C₂H₄ line list of the *HITRAN* database. Consequently,
 2 Eq. (22) has been used for all the levels to calculate the statistical weights g_{rot} .

3 6.3.39. CH₃OH (molecule 39)

4 The statistical weights g_s are 8 for the *A* rotational levels if the splitting is not observable (4 for
 5 each member of a resolvable *A*± doublet) and 4 for the *E* rotational levels [39].

6 No hyperfine structure is reported in the CH₃OH line list of the *HITRAN* database, and Eq.
 7 (22) has been used for all the levels to calculate the statistical weights g_{rot} .

11 7. Relation between the Einstein *A*-coefficient, the weighted square of the transition moment, the line 12 intensity and the oscillator strength

13 The relation between the Einstein *A*-coefficient and the unweighted square of the transition
 14 moment depends on the type of a transition [12,21]

15
 16 For an electric-dipole transition $g_2 A_{21} = \frac{16\pi^3}{3h\epsilon_0} v^3 |R_{12}|^2$. (32)

17
 18 For a magnetic-dipole transition $g_2 A_{21} = \frac{16\pi^3 \mu_0}{3h} v^3 |R_{12}|^2$. (33)

19
 20 For an electric-quadrupole transition $g_2 A_{21} = \frac{8\pi^5}{5h\epsilon_0} v^5 |R_{12}|^2$. (34)

21
 22 The factors ϵ_0 and μ_0 are the fundamental physical constants, the permittivity and the
 23 permeability of vacuum, respectively. The transition moment squared has the units C² m², A² m⁴
 24 and C² m⁴ in Eqs. (32), (33) and (34), respectively. Eq. (32) can be found in Refs. [6,13].

25 From Eqs. (20) and (32) in cgs units (for conversion to cgs electrostatic units, ϵ_0 should be
 26 replaced by $1/4\pi$ [12]), we obtain the relation linking the line intensity S_{HIT} (in cm⁻¹ (molecule
 27 cm⁻²)⁻¹) to the weighted square of the transition moment $\mathfrak{R}_{12} \equiv 1/g_1 |R_{12}|^2$ (in Debye²,
 28 1 Debye = 10⁻¹⁸ cgsesu), which has been used for the previous *HITRAN* compilations
 29 [2,5,6,10,15] (the majority of the transitions in *HITRAN* are electric-dipole)

30
 31
 32 $S_{\text{HIT}} = I_a \frac{v_0}{Q_{\text{tot}}(T_0)} \frac{8\pi^3}{3hc} e^{-c_2 E''/T_0} (1 - e^{-c_2 v_0/T_0}) g_1 \mathfrak{R}_{12} \times 10^{-36}$. (35)

33
 34 In many studies, especially in astrophysics, we find the dimensionless oscillator strength, also
 35 called the *f*-value, instead of the Einstein *A*-coefficient, the transition moment or the line intensity.
 36 It is thus useful to recall the relation between the oscillator strength and the Einstein *A*-coefficient
 37 [6,13,14]

38
 39 $f_{12} = \frac{g_2}{g_1} \frac{\epsilon_0 m c}{2\pi e^2 v_0^2} A_{21}$, (36)

40
 41 where *e* and *m* are the charge and the mass of the electron, respectively. One can note that $g_1 f_{12}$ is
 42 also called the weighted oscillator strength. The relation between the oscillator strength and the

line intensity or the transition moment squared can easily be obtained by combining Eq. (36) with Eq. (20) or Eqs. (32)–(34), respectively, depending on the type of the transition.

8. Non-additivity of the Einstein A -coefficients

Contrary to the line intensities, the Einstein A -coefficients and the oscillator strengths are not additive. If we consider the upper state noted η' and the lower state noted η'' with the degeneracies $d_{\eta'}$ and $d_{\eta''}$, respectively, the Einstein A -coefficient $A_{\eta' \rightarrow \eta''}$ for spontaneous emission from the state η' to η'' can be expressed from the Einstein A -coefficients $A_{\eta' \xi' \rightarrow \eta'' \xi''}$ for spontaneous emissions from the substates $\eta' \xi'$ to $\eta'' \xi''$ using the following equation [6]

$$A_{\eta' \rightarrow \eta''} = \frac{1}{d_{\eta'}} \sum_{\substack{\xi', \xi'' \\ \xi', \xi''}} A_{\eta' \xi' \rightarrow \eta'' \xi''}. \quad (37)$$

For the lines with the hyperfine structure listed in *HITRAN*, we can express the Einstein A -coefficient of the transition from the Einstein A -coefficients of its hyperfine components

$$A_{e'v'J' \rightarrow e''v''J''} = \frac{1}{d_{J'}} \sum_{F_1, F_1'} A_{e'v'J'F_1 \rightarrow e''v''J''F_1'}, \quad (38)$$

where the indices e, v, J , and F_1 (denoted F in *HITRAN*, see Section 6.2) correspond to the electronic, vibrational, rotational, and hyperfine levels, respectively. In this case, $d_{J'}$ represents the hyperfine multiplicity of the level J' , that is, the number of its hyperfine sublevels F_1' .

9. Comparisons between the Einstein A -values from this work and submillimeter catalogs

It is informative to make comparisons of the values calculated here with similar quantities used in archival databases. Verdes et al. [40] made extensive comparisons for the partition sum data between *HITRAN* and the JPL submillimeter catalog [25]. The JPL catalog is limited to lines below 300 cm^{-1} , and is focused on constituents in the interstellar medium. Verdes et al. [40] concluded that the agreement between the *HITRAN* and *JPL* values was generally quite good, provided that one took into account that the JPL catalog did not contain the vibrational partition sum. Some differences that did exist, for example for nitric oxide, have since been corrected in the **HITRAN** calculation.

We made a comparison of Einstein A -coefficients between *HITRAN* [1] and *CDMS* [3]. Similar to the JPL catalog, *CDMS* is limited to the submillimeter region, and isotopologues in common between *HITRAN* and *CDMS* are not numerous. However, the *CDMS* web-site allows users to have both the intensities and Einstein A -coefficients, as does the new format of *HITRAN* [1]. For example, Fig. 2 represents the ratio of the Einstein A -coefficients from Refs. [3,1] for the (0–0) bands of $^{12}\text{C}^{16}\text{O}$ and $^{12}\text{C}^{18}\text{O}$ isotopologues. These plots show systematic discrepancies of about 1–3% increasing with the wavenumber. The two databases also give very similar results for the (01¹0–01¹0) band of $\text{H}^{12}\text{C}^{14}$ (the differences are less than 1%, see Fig. 3). Comparisons are also possible for the pure-rotational bands of H_2CO , and SO_2 . The difference between *HITRAN* and *CDMS* values does not exceed 2% for $\text{H}_2^{13}\text{C}^{16}\text{O}$ and $\text{H}_2^{12}\text{C}^{18}\text{O}$; however the discrepancies reach




Fig. 2. Comparisons of the Einstein A -coefficients for the (0-0) bands of the $^{12}\text{C}^{16}\text{O}$ and $^{12}\text{C}^{18}\text{O}$ isotopologues.




Fig. 3. Comparison of the Einstein A -coefficients for the (01¹₀-01¹₀) band of the $\text{H}^{12}\text{C}^{14}\text{N}$ isotopologue.

1 almost 25% for $\text{H}_2^{12}\text{C}^{16}\text{O}$ and $^{32}\text{S}^{16}\text{O}_2$. In these cases the ratios of the Einstein A -coefficients are
 2 almost ($\pm 5\%$) identical with the ratios of the line intensities. The discrepancies are probably
 3 attributable to the different values of the dipole-moment expansion used in the two databases.

5 Acknowledgements

7 The authors thank V. Boudon, I. Kleiner, A. Perrin, and J.B. Tatum for useful discussions and
 9 comments concerning the calculation of statistical weights. We also wish to acknowledge P.
 10 Bernath and J.-Y. Mandin for helpful comments regarding the manuscript.

11 The current effort has been supported by the NASA Earth Observing System (EOS), Grant
 12 NAG5-13534, and funding from the Office of the Secretary of Defense (OSD) High Energy Laser
 13 Joint Technology Office. One of the authors (RRG) is pleased to acknowledge support of this
 14 research by the National Science Foundation (NSF) through Grant No. ATM-0242537.

17 References

- 19 [1] Rothman LS, Jacquemart D, Barbe A, Benner DC, Birk M, Brown LR, et al. The HITRAN 2004 molecular
 20 spectroscopic database. *JQSRT* 2005;96:139–204.
- 21 [2] Rothman LS, Barbe A, Benner DC, Brown LR, Camy-Peyret C, Carleer MR, et al. The HITRAN molecular
 22 spectroscopic database: edition of 2000 including updates through 2001. *JQSRT* 2003;82:5–44.
- 23 [3] Müller HSP, Thorwirth S, Roth DA, Winnewisser G. The Cologne database for molecular spectroscopy, CDMS.
 24 *Astron Astrophys* 2001;370:L49–52.
- 25 [4] Goldman A, Gamache RR, Perrin A, Flaud JM, Rinsland CP, Rothman LS. HITRAN partition function and
 26 weighted transition-moments squared. *JQSRT* 2000;66:455–86.
- 27 [5] Rothman LS, Rinsland CP, Goldman A, Massie ST, Edwards DP, Flaud JM, et al. The HITRAN molecular
 28 spectroscopic database and HAWKS (HITRAN atmospheric workstation): 1996 edition. *JQSRT*
 29 1998;60:665–710.
- 30 [6] Gamache RR, Rothman LS. Extension of the HITRAN database to non-LTE applications. *JQSRT*
 31 1992;48:519–25.
- 32 [7] Rothman LS, Hawkins RL, Wattson RB, Gamache RR. Energy levels, intensities, and linewidths of atmospheric
 33 carbon dioxide bands. *JQSRT* 1992;48:537–66.
- 34 [8] Jacquemart D, Claveau C, Mandin JY, Dana V. Line intensities of hot bands in the $13.6\ \mu\text{m}$ spectral region of
 35 acetylene $^{12}\text{C}_2\text{H}_2$. *JQSRT* 2001;69:81–101.
- 36 [9] Weber M, Blass WE, Halsey GW, Hillman JJ. ℓ -Resonance perturbation of IR intensities in $^{12}\text{C}_2\text{H}_2$ near $13.7\ \mu\text{m}$.
 37 *J Mol Spectrosc* 1994;165:107–23.
- 38 [10] Rothman LS, Gamache RR, Tipping RH, Rinsland CP, Smith MAH, Benner DC, et al. The HITRAN molecular
 39 database: editions of 1991 and 1992. *JQSRT* 1992;48:469–507.
- 40 [11] Teffo JL, Lyulin OM, Perevalov VI, Lobodenko EI. Application of the effective operator approach to the
 41 calculation of $^{12}\text{C}^{16}\text{O}_2$ line intensities. *J Mol Spectrosc* 1998;187:28–41.
- 42 [12] Tatum JB. The interpretation of intensities in diatomic molecular spectra. *Astrophys J Suppl Ser* 1967;14:21–56;
 43 Tatum JB. Erratum: interpretation of intensities in diatomic molecular spectra. *Astrophys J Suppl Ser* 1971;22:388.
- [13] Hilborn RC. Einstein coefficients, cross sections, f values, dipole moments, and all that. *Am J Phys* 1982;50:982–6.
- [14] Penner SS. Quantitative molecular spectroscopy and gas emissivities. Reading, MA: Addison-Wesley; 1959.
- [15] Rothman LS, Gamache RR, Goldman A, Brown LR, Toth RA, Pickett HM, et al. The HITRAN database: 1986
 edition. *Appl Opt* 1987;26:4058–97.
- [16] Pugh LA, Rao KN. Intensities from infrared spectra. *Mol Spectrosc Mod Res* 1976;II:165–227.

- 1 [17] Mohr PJ, Taylor BN. CODATA recommended values of the fundamental physical constants: 1998. *Rev Mod Phys* 2000;72:351–495.
- 3 [18] Fischer J, Gamache RR, Goldman A, Rothman LS, Perrin A. Total internal partitions sums for molecular species on the 2000 edition of the HITRAN database. *JQSRT* 2003;82:401–12.
- 5 [19] Gamache RR, Kennedy S, Hawkins RL, Rothman LS. Total internal partition sums for molecules in the terrestrial atmosphere. *J Mol Struct* 2000;517–8:407–25.
- 7 [20] Fischer J, Gamache RR. Partition sums for non-local thermodynamic equilibrium applications. *JQSRT* 2002;74:273–84.
- 9 [21] Tatum JB. Structure and intensities of microwave lines in the spectra of diatomic molecules. *Astrophys J Suppl Ser* 1986;60:433–74.
- 11 [22] Gamache RR, Goldman A. Einstein A coefficient, integrated band intensity, and population factors: application to the $a^1\Delta_g - X^3\Sigma_g^-(0,0)$ O₂ band. *JQSRT* 2001;69:389–401.
- 13 [23] Gamache RR, Goldman A, Rothman LS. Improved spectral parameters for the three most abundant isotopomers of the oxygen molecule. *JQSRT* 1998;59:495–509.
- 15 [24] Herzberg G. *Molecular spectra and molecular structure, II. Infrared and Raman spectra of polyatomic molecules.* Princeton: D Van Nostrand; 1945.
- 17 [25] Pickett HM, Poynter RL, Cohen EA, Delitsky ML, Pearson JC, Müller HSP. Submillimeter, millimeter, and microwave spectral line catalog. *JQSRT* 1998;60:883–90.
- 19 [26] Bunker PR, Jensen P. *Molecular symmetry and spectroscopy.* Ottawa: NRC Research Press; 1998.
- 21 [27] Herzberg G. *Molecular spectra and molecular structure, I. Spectra of diatomic molecules.* Princeton: D Van Nostrand; 1950.
- 23 [28] Papoušek D, Aliev MR. *Molecular vibrational/rotational spectra.* Prague: Academia; 1982.
- 25 [29] Hougen JT. Classification of rotational energy levels for symmetric-top molecules. *J Chem Phys* 1962;37:1433–41.
- 27 [30] Chackerian Jr C, Brown LR, Lacombe N, Tarrago G. Methyl chloride ν_5 region line shape parameters and rotational constants for the ν_2 , ν_5 and $2\nu_3$ vibrational bands. *J Mol Spectrosc* 1998;191:148–57.
- 29 [31] Jacquemart D, Mandin JY, Dana V, Claveau C, Vander Auwera J, Herman M, et al. The IR acetylene spectrum in HITRAN: update and new results. *JQSRT* 2003;82:363–82.
- 31 [32] Ruoff A. From special unitary groups SU(2I + 1) to nuclear spin statistical weights. *J Mol Spectrosc* 2003;219:177–85.
- 33 [33] Hougen JT. Perturbations in the vibration–rotation–torsion energy levels of an ethane molecule exhibiting internal rotation splittings. *J Mol Spectrosc* 1980;82:92–116.
- 35 [34] Berger H. Classification of energy levels for polyatomic molecules. *J Phys* 1977;38:1371–5.
- 37 [35] Galavis ME, Mendoza C, Zeippen CJ. Atomic data from the IRON Project. XXII. Radiative rates for forbidden transitions within the ground configuration of ions in the carbon and oxygen isoelectronic sequences. *Astron Astrophys Suppl* 1997;123:159–71.
- [36] Vastel C, Caux E, Ceccarulli C, Castets A, Gry C, Baluteau JP. Large [O]/[CO] ratios in cold molecular clouds towards W 49N. *Astron Astrophys* 2000;357:994–1000.
- [37] Oomens J, Reuss J, Mellau GCh, Klee S, Gulaczyk I, Fayt A. The ethylene hot band spectrum near 3000 cm⁻¹. *J Mol Spectrosc* 1996;180:236–48.
- [38] Bach M, Georges R, Herman M, Perrin A. Investigation of the fine structure in overtone absorption bands of ¹²C₂H₄. *Mol Phys* 1999;97:265–77.
- [39] Xu LH, Lees RM, Wang P, Brown LR, Kleiner I, Johns JWC. New assignments, line intensities, and HITRAN database for CH₃OH at 10 μm. *J Mol Spectrosc* 2004;228:453–70.
- [40] Verdes CL, von Engeln A, Buehler SA, Perrin A. Partition function data and impact on retrieval quality for an mm/sub-mm limb sounder. *JQSRT* 2005;90:217–38.