

The method of undetermined coefficients can be used to find a particular solution y_p of a nonhomogeneous linear d.e. **if** the d.e. has constant coefficients **and** the nonhomogeneous term is a polynomial, an exponential, a sine or a cosine, or a sum or product of these.

If the d.e. has variable coefficients and/or the nonhomogeneous term is something other than a polynomial, exponential, sine, cosine, or sum or product of these, you must use another method (e.g. variation of parameters).

1. If the nonhomogeneous term is a polynomial, try guessing a polynomial of the same degree for y_p .

Example. $y'' - 3y' + 2y = 4x$

Nonhomogeneous term: $4x$ (polynomial of degree 1).

Guess: $y_p = Ax + B$ (a polynomial of degree 1).

Note: You need the constant term B in your guess for y_p even though there is no constant in the nonhomogeneous term.

2. If the nonhomogeneous term is an exponential function, try guessing an exponential function of the same form for y_p .

Example. $y'' - 3y' + 2y = 6e^{-x}$

Nonhomogeneous term: $6e^{-x}$ (an exponential function).

Guess: $y_p = Ae^{-x}$ (an exponential function of the same form).

3. If the nonhomogeneous term is a sine or a cosine, try guessing a combination of sine and cosine of the same angular frequency for y_p .

Example. $y'' - 3y' + 2y = 10 \sin(2x)$

Nonhomogeneous term: $10 \sin(2x)$ (sine function with angular frequency 2).

Guess: $y_p = A \sin(2x) + B \cos(2x)$ (combination of sine and cosine w. angular frequency 2).

Note: You need the cosine term $B \cos(2x)$ in your guess for y_p even though there is no cosine in the nonhomogeneous term.

4. If the nonhomogeneous term is a sum of a polynomial, an exponential function, and/or a sine or cosine, try guessing a sum of these functions for y_p .

Example. $y'' - 3y' + 2y = 4x + 10 \sin(2x)$

Nonhomogeneous term: $4x + 10 \sin(2x)$ (sum of a polynomial of degree 1 and a sine with angular frequency 2.)

Guess: $y_p = Ax + B + C \sin(2x) + D \cos(2x)$

5. If the nonhomogeneous term is a product of a polynomial, an exponential function, and/or a sine or cosine, try guessing a product of these functions for y_p .

Example. $y'' - 3y' + 2y = 36xe^{-x}$

Nonhomogeneous term: $36xe^{-x}$ (product of a polynomial of degree 1 and an exponential.)

Guess: $y_p = (Ax + B)e^{-x}$

Note: You need the Be^{-x} term in your guess for y_p even though there is no such term in the nonhomogeneous term.

OVER

IMPORTANT EXCEPTION TO THE ABOVE RULES

If any part of your guess for y_p is a part of the complementary solution y_c , you must multiply that part of your y_p guess by x .

Example 1. $y'' - 3y' + 2y = e^x$

The homogeneous equation is $y'' - 3y' + 2y = 0$, which has characteristic equation $r^2 - 3r + 2 = 0$.

The roots of the characteristic equation are 1 and 2, so the complementary solution is $y_c = c_1e^x + c_2e^{2x}$.

Nonhomogeneous term: e^x

Usual guess: $y_p = Ae^x$. A term of this form already appears in y_c , so we must multiply our guess by x .

Correct guess: $y_p = Axe^x$.

Example 2. $y'' - 2y' + y = e^x$

The homogeneous equation is $y'' - 2y' + y = 0$, which has characteristic equation $r^2 - 2r + 1 = 0$.

The root of the characteristic equation is 1 (with multiplicity 2), so the complementary solution is $y_c = c_1e^x + c_2xe^x$.

Nonhomogeneous term: e^x .

Usual guess: $y_p = Ae^x$. A term of this form already appears in y_c , so we multiply our guess by x : $y_p = Axe^x$. This new guess is still of the same form as part of y_c , so we must multiply by x again.

Correct guess: $y_p = Ax^2e^x$.