

Math 491, Problem Set #7: Solutions

1. One basis for the space of polynomials of degree less than d is the monomial basis $1, t, t^2, \dots, t^{d-1}$. Another is the shifted monomial basis $1, (t+1), (t+1)^2, \dots, (t+1)^{d-1}$. Call these bases u_1, \dots, u_d and v_1, \dots, v_d respectively.

- (a) Derive a formula for the entries of the change-of-basis matrix M expressing the u_i 's as linear combinations of the v_j 's.

We seek a d -by- d matrix M that, when multiplied on the right by the column vector e_i (with a 1 in the i th position and a 0 everywhere else), gives a column vector $(c_1, c_2, \dots, c_d)^T$ such that $u_i = c_1v_1 + c_2v_2 + \dots + c_dv_d$. Now $u_i = t^{i-1} = ((t+1) - 1)^{i-1} = \sum_{j=0}^{i-1} \binom{i-1}{j} (t+1)^j (-1)^{i-1-j} = \sum_{j=0}^{i-1} \binom{i-1}{j} v_{j+1} (-1)^{i-1-j} = \sum_{j=1}^i \binom{i-1}{j-1} v_j (-1)^{i-j}$, so $c_j = (-1)^{i-j} \binom{i-1}{j-1}$ (which gets interpreted as 0 for $j > i$). Hence

$$M_{j,i} = \begin{cases} (-1)^{i-j} \binom{i-1}{j-1} & \text{for } 1 \leq j \leq i \leq n, \\ 0 & \text{otherwise.} \end{cases}$$

(Note: I didn't specify whether the vectors were to be treated as row-vectors or column-vectors, or equivalently, whether the change-of-basis matrix was supposed to be applied on the right or on the left. If you adopted the row-vector approach, you would find that the answers you got for parts (a) and (b) are reversed, relative to mine.)

- (b) Derive a formula for the entries of the change-of-basis matrix N expressing the v_j 's as linear combinations of the u_i 's.

This one is even easier: $v_j = (t+1)^{j-1} = \sum_{i=0}^{j-1} \binom{j-1}{i} t^i = \sum_{i=1}^j \binom{j-1}{i-1} u_i$
so

$$N_{i,j} = \begin{cases} \binom{j-1}{i-1} & \text{for } 1 \leq i \leq j \leq n, \\ 0 & \text{otherwise.} \end{cases}$$

- (c) From the description of M and N as basis-change matrices, we know that $MN = NM = I$. Forgetting for the moment what M

and N mean, rewrite the assertions $MN = NM = I$ as binomial coefficient identities, and prove them either algebraically or bijectively.

The assertion $MN = I$ can be rewritten as $\sum_j M_{i,j} N_{j,k} = \delta_{i,k}$, where $\delta_{i,j}$ is 1 if $i = j$ and 0 otherwise. That is, $\sum (-1)^{j-i} \binom{j-1}{i-1} \binom{k-1}{j-1} = \delta(i, k)$ where the sum is over all j such that $i \leq j \leq k$. For convenience, we shift indices and write this as

$$\sum (-1)^{j-i} \binom{j}{i} \binom{k}{j} = \delta(i, k)$$

where the sum is still over all j such that $i \leq j \leq k$.

Algebraic proof: The sum in question is the coefficient of x^{k-i} in the product of $\binom{i}{i} - \binom{i+1}{i}x + \binom{i+2}{i}x^2 - \dots + (-1)^{k-i} \binom{k}{i}x^{k-i} + \dots$ and $\binom{k}{k} + \binom{k}{k-1}x + \binom{k}{k-2}x^2 + \dots + \binom{k}{i}x^{k-i} + \dots + \binom{k}{0}x^k$. The first factor can be recognized as $(1+x)^{-(i+1)}$ (by the binomial theorem) and the latter can be recognized as $(1+x)^k$. So the product is $(1+x)^{k-i-1}$. The coefficient of x^{k-i} in the formal power series expansion of $(1+x)^{k-i-1}$ is 0 as long as $k-i-1$ is non-negative, since in that case $(1+x)^{k-i-1}$ is just a polynomial of degree less than $k-i$. However, when $i = k$, $(1+x)^{k-i-1}$ becomes the formal power series $(1+x)^{-1} = 1 - x + x^2 - x^3 + \dots$, in which the coefficient of x^{k-i} is just the constant term 1.

Combinatorial proof: Given a set C of size k , $\sum (-1)^{j-i} \binom{j}{i} \binom{k}{j}$ counts the number of ways to choose a subset $B \subset C$ of size j and a subset $A \subset B$ of size i , where a choice of A, B, C counts as positive or negative according to whether the number of elements of B that are not in C is even or odd. If we hold the subset A fixed and do a signed enumeration of the sets B satisfying $A \subset B \subset C$, we find that the signed count is 1 if $A = C$ and 0 otherwise. (Reason: This is just like signed enumeration of the subsets of $C \setminus A$, where a set counts as positive or negative according to whether it has an even or odd number of elements.) If $i = k$, there is exactly one set A , namely C itself, whose aggregate contribution is non-zero, and in this case the aggregate contribution is 1; whereas if $i < k$, all the aggregate contributions vanish. This proves the identity.

The assertion $NM = I$ can be rewritten as $\sum_j N_{i,j}M_{j,k} = \delta_{i,k}$, that is, $\binom{j-1}{i-1}(-1)^{k-j}\binom{k-1}{j-1} = \delta_{i,j}$. Re-indexing, we write $(-1)^{k-j}\binom{j}{i}\binom{k}{j} = \delta_{i,j}$. The proofs are similar to what appeared above.