

ENVI.2030L – Rock Identification

I. Introduction

The bulk of the earth's crust is composed of relatively few minerals. These can be mixed together, however, to give an endless variety of rocks - aggregates of minerals, rock fragments and glass. All rocks are classified on the basis of their mineral content (or other components if minerals are not present) and texture. The system of classification and the textural terminology are different, however, for the three groups (igneous, metamorphic, sedimentary) of rocks. It is therefore important to determine the rock group before attempting to classify a rock.

II. Classification into rock groups

Rocks can be assigned to one of the three groups by considering the following characteristics of each rock group:

Igneous. Igneous rocks have solidified from molten matter (magma) and are composed of individual, interlocking crystals. If solidification occurred very rapidly the rock may be glassy, contain glass shards, or be filled with holes (vesicles) which are formed when gas escapes from the magma.

Sedimentary. Sedimentary rocks are composed of particles derived from pre-existing rocks, or materials precipitated from solution, or organic matter. In the field sedimentary rocks show distinct layering which may not be apparent in small hand specimens. Sedimentary rocks are composed of individual grains cemented together. Fossils are almost exclusively restricted to sedimentary rocks.

Metamorphic. Metamorphic rocks are derived from previously existing rocks in response to changes in pressure, temperature and fluid content. These process change the mineral composition and/or texture of the pre-existing rock. In general, metamorphic rocks have a banded or foliated appearance (foliated - appear to be composed of thin layers pressed together). If the rock is composed of one mineral it may have a blocky appearance and be composed of large crystals that are densely intergrown.

III. Igneous rocks

Igneous rocks are classified on the basis of their texture and mineral composition.

Textural terms:


1. Pegmatitic - grains greater than 1.2 cm in diameter.
2. Phaneritic - grains can be seen with the unaided eye.
3. Aphanitic - grains cannot be seen with the unaided eye (i.e. you need a hand lens or microscope).
4. Porphyritic-phaneritic - two distinct grain sizes. Smaller grains can be seen with the unaided eye.
5. Porphyritic-aphanitic - two distinct grain sizes. Smaller grains cannot be seen with the unaided eye.
6. Amygdaloidal - vesicles in aphanitic rock filled with a mineral.
7. Glassy - rock looks like a glass. No crystalline structure. Example - obsidian.

8. Scoriaceous - many holes (vesicles) found in the rock. Frothy appearance. Examples – pumice (usually light colored), scoria (usually red or black in color).
9. Fragmental - rock composed of angular fragments. Example - igneous breccia

Identification of common minerals in igneous rocks:

1. Quartz - smokey appearance, vitreous luster, hardness of 7, no cleavage.
2. Orthoclase feldspar - light in color, two right angle cleavages, hardness of 6.
3. Plagioclase feldspar - light to dark in color, two right angle cleavages, hardness of 6.
4. Muscovite - cleavage results in flakes, light in color.
5. Biotite - cleavage results in flakes, dark in color.
6. Hornblende - dark green in color, splintery, elongate, 60° - 120° cleavage angle.
7. Augite - black in color, hard, poor cleavage, 90° - 90° cleavage.
8. Olivine - green in color, granular appearance.

A generalized classification scheme for the igneous rocks is given in Figure 1. Note that scoria does not appear in Figure 1, but you may want to use this rock name when you classify your unknowns. **Scoria** is a highly vesicular, dark colored (brown, black, or red) volcanic rock that may or may not contain crystals (phenocrysts).



© 2010 Tasa Graphic Arts, Inc.

Figure 1. Igneous rock classification scheme

IV. Sedimentary rocks

Sedimentary rocks can be broadly divided into three groups:

1. Clastic - formed from the accumulation of mineral and rock fragments.
 - Breccia - large angular fragments
 - Conglomerate - large rounded fragments, pebbles
 - Sandstone - composed of sand sized particles
 - Shale - composed of mud sized particles, has fissility, a dull thud when struck
 - Tillite - composed of all sizes of particles from silt to large fragments, glacial origin

2. Organic and biochemical - rocks produced from organic remains or minerals formed by organisms.
 - Limestone - composed of calcite or aragonite
 - Lithographic - extremely fine grained
 - Fossiliferous - contains shell fragments
 - Coquina - poorly cemented shell fragments
 - Chalk - accumulation of skeletons of one-celled animals that secrete calcareous shells
 - Dolomite - composed of dolomite
 - Siliceous - composed of silica
 - Flint - black in color, hard
 - Chert - light in color, hard
 - Carbonaceous - composed predominately of organic material
 - Peat - loose and porous accumulation of organic matter
 - Lignite - a brown coal with a fibrous woody structure
 - Bituminous coal - soft coal, dull black in color
 - Anthracite coal - hard coal, shiny black in color
 - Bog iron ore - composed of hydrated iron oxide (goethite)

3. Chemical precipitates - rocks formed by precipitation of crystalline material from solution. Examples are gypsum and rock salt.

A generalized classification scheme for the sedimentary rocks is given in Figure 2.

V. Metamorphic rocks

Metamorphic rocks are divided into two groups based on the presence or absence of foliation. Foliation varies from very fine to coarse foliation (mica flakes clearly visible) to gneissic textures in which the minerals are segregated into layers. A simplified classification scheme for the metamorphic rocks is given in Figure 3.

SEDIMENTARY ROCK ANALYSIS AND CLASSIFICATION						
STEP 1: What is the rock's composition?		STEP 2: What are the rock's textural and other distinctive properties?		STEP 3: Rock Name(s)		
DETRITAL (CLASTIC)	Mainly rock fragments or mineral grains (quartz, feldspar, clay) weathered from other rocks	Mainly gravel (≥ 2 mm)	Rounded grains	CONGLOMERATE		
			Angular grains	BRECCIA		
		Mainly sand (1/16 – 2 mm)	Mostly quartz grains	QUARTZ SANDSTONE	SANDSTONE	
			Mainly feldspar and quartz	ARKOSE		
			Sand is mixed with much silt and/or clay (mud)	GRAYWACKE		
		Mainly Mud (< 1/16 mm)	Mostly silt (1/256 – 1/16 mm)	Breaks into blocks or layers	SILTSTONE	MUDSTONE
				Crumbles or breaks into blocks	CLAYSTONE	
Mostly clay (< 1/256 mm)	Fissile (splits easily)		SHALE			
BIOCHEMICAL: Mainly fossil shells or plant fragments	Mainly plant fragments or charcoal	Dull brown with visible plant fragments	Porous and easy to break apart the plant fragments	PEAT		
		Black	Dense and brittle or porous and sooty	BITUMINOUS COAL		
BIOCHEMICAL: Mainly fossil shells or plant fragments	Mainly fossil shells, shell fragments, or microfossils Effervesces in dilute HCl	Mostly visible shells and shell fragments cemented into a dense mass		CALCIRUDITE		
		Mostly sand-sized fragments. May have a few larger shells.		CALCARENITE		
		Mostly very fine grained to microcrystalline mass of calcite and microfossils		MICRITE		
		Porous, poorly cemented mass of shells and shell fragments		COQUINA		
		Mostly very fine grained, earthy, chalky, light-colored mass of microfossils		CHALK		
		Crystalline to microcrystalline bands of calcite crystals		TRAVERTINE		
CHEMICAL (INORGANIC): Chemically precipitated crystals	Mainly crystals of calcite or aragonite, CaCO_3 Effervesces in dilute HCl	Spherical grains like tiny beads (< 2 mm) with concentric laminations		OOLITIC LIMESTONE		
		Microcrystalline		DOLOSTONE		
	Mainly dolomite $\text{CaMg}(\text{CO}_3)_2$	Microcrystalline, conchoidal fracture		CHERT		
		Crystals formed as inorganic chemical precipitates		ROCK SALT		
	Mainly varieties of quartz, SiO_2 (chalcedony, flint, chert, opal, jasper, etc.)	Scratches glass		CHERT		
	Mainly halite, NaCl	Salty taste		ROCK SALT		
Mainly gypsum, $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$	Crystals formed as inorganic chemical precipitates		ROCK GYPSUM			
Mostly iron-bearing minerals, like limonite and hematite	Amorphous or microcrystalline		IRONSTONE			

Figure 2. Sedimentary rock classification scheme.

Rock Name	Texture	Grain Size	Comments	Parent Rock
Slate	Foliated	Very fine	Excellent rock cleavage, smooth dull surfaces	Shale, mudstone, or siltstone
Phyllite		Fine	Breaks along wavy surfaces, glossy sheen	Slate
Schist		Medium to Coarse	Micas dominate, scaly foliation	Phyllite
Gneiss		Medium to Coarse	Compositional banding due to segregation of minerals	Schist, granite, or volcanic rocks
Marble	Nonfoliated	Medium to coarse	Interlocking calcite or dolomite grains	Limestone, dolostone
Quartzite		Medium to coarse	Fused quartz grains, massive, very hard	Quartz sandstone
Anthracite		Fine	Shiny black organic rock that may exhibit conchoidal fracture	Bituminous coal

Copyright © 2006 Pearson Prentice Hall, Inc.

Figure 3. Classification scheme for the metamorphic rocks

VI. Classification of unknown rock specimens

Separate your unknowns into igneous, sedimentary and metamorphic groups. The instructor will check your groupings. Identify the rocks within each group using texture and mineralogy. Enter the rock name and its characteristics on the data sheet (next page).

Data Table for Rock Identification Exercise

Rock Name	Texture	Components
Igneous Rocks		
Metamorphic Rocks		
Sedimentary Rocks		