

# Non-specific Interaction

- # Electrostatic in nature
- # Limits effectiveness of ion in solution
- # Use concept of **activity** to quantify effect

(effective concentration)

$$a_i = [i]_F \gamma_F(i)$$

where  $a_i$  = activity of ion  $i$

$[i]_F$  = free ion conc. (m)

$\gamma_F(i)$  = activity coefficient

of ion  $i$

In short

$$a = [i] \gamma$$

# Activity of Individual Ion Influenced by Other Ions

## # Ionic Strength of solution

$$I = 0.5 \sum Z^2 m$$

where I = ionic strength

Z = charge on ion

m = molal conc.

(molarity or molinity

can also be used)

$$a = [i] \gamma$$

**Table 4.1** *Concentrations of the major constituents in surface seawater**At salinity (PSS 1978): S = 35.000*

	<i>g/kg</i>	<i>mmol/kg</i>	<i>mM</i>
Na <sup>+</sup>	10.781	468.96	480.57
K <sup>+</sup>	0.399	10.21	10.46
Mg <sup>++</sup>	1.284	52.83	54.14
Ca <sup>++</sup>	0.4119	10.28	10.53
Sr <sup>++</sup>	0.00794	0.0906	0.0928
Cl <sup>-</sup>	19.353	545.88	559.40
SO <sub>4</sub> <sup>-</sup>	2.712	28.23	28.93
HCO <sub>3</sub> <sup>-</sup>	0.126	2.06	2.11
Br <sup>-</sup>	0.0673	0.844	0.865
B(OH) <sub>3</sub>	0.0257	0.416	0.426
F <sup>-</sup>	0.00130	0.068	0.070

SW Density = 1.024763 kg/L at 20 °C (Pilson 1998)

# Major Components of SW

- #  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Cl}^-$  and  $\text{SO}_4^{2-}$  are most abundant
- # Account for 98.5 % of dissolved species in SW
- # Have major influence on SW density
- # Have long residence time in the ocean
- # Generally exhibit conservative behavior
  - Concentration influenced by physical processes such as evaporation & precipitation, not chemical or biological processes
- # Discussing completely dissolved species

# Element Concentrations in Average River & Average Ocean Water with Residence Times

	Conc. Mean River ( $10^{-6}$ moles/kg)	Conc. Mean Sea ( $10^{-6}$ moles/kg)	$\tau$ (yrs)
Na	$2.2 \times 10^2$	$4.7 \times 10^5$	$8.3 \times 10^7$
Mg	$1.6 \times 10^2$	$5.3 \times 10^4$	$1.3 \times 10^7$
Al	1.9	( $3 \times 10^{-2}$ )	$6.2 \times 10^2$
Si	$1.9 \times 10^2$	$1.0 \times 10^2$	$2.0 \times 10^4$
P	1.3	2.3	$6.9 \times 10^4$
S	-	$2.8 \times 10^4$	-
Cl	-	$5.5 \times 10^5$	-
Ar	-	$1.5 \times 10^1$	-
K	$3.4 \times 10^1$	$10.2 \times 10^3$	$1.2 \times 10^7$
Ca	$3.6 \times 10^2$	$10.3 \times 10^3$	$1.1 \times 10^6$

# Cycling of SW Components

“The sea is a way station for the products of continental erosion. All substances received by the sea are ultimately passed along to the sediment...tectonic forces...eventually push the material buried in this way back above sea level where it becomes subject to erosion. Then another trip through the sea begins.”

Broecker and Peng (1982)

# Cycling of SW Components

- # Most components are recycled many times within SW by a variety of processes
- # Can determine residence times ( $\tau$ ) in ocean
- # Constituents can be classified as:
  - Biolimiting – totally depleted in surface water
  - Biointermediate – partially depleted
  - Biounlimited – no measurable depletion
  - Noncycling – reactive & removed

Broecker and Peng (1982)

# SW Composition


The composition of SW generally reflects two factors:

- 1) The relative abundance of the substance in river water (i.e., the input)
- 2) The presence of removal mechanisms that result in entrapment of the material in sediments (i.e., the output)


# Major Components of SW

- #  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Cl}^-$  and  $\text{SO}_4^{2-}$  are most abundant
- # Account for 98.5 % of dissolved species in SW
- # Have major influence on SW density
- # Have long residence time in the ocean
- # Generally exhibit **conservative** behavior
  - Concentration influenced only by physical processes such as evaporation & precipitation, not chemical or biological processes
- # Discussing completely dissolved species

# Conservative Mixing



**Ion Concentration**



**Volume Ratio of Water Mass**

# Marcet Principle (1819)

- # Relative composition of sea salt is nearly the same worldwide, i.e., major constituents are conservative
- # Constancy of Composition
- # Principle of Constant Composition (Pilson)
- # Rule of Constant Proportions (Libes)
- # First Law of Chemical Oceanography (Kester)
- # Several exceptions to the rule

# Exceptions to the Rule

(or non-conservative behavior)

- # Caused by processes such as:  
Reduction, Dissolution, Evaporation, etc.
- # Estuaries & Marginal Seas – largely input of river water of different composition & other processes also (e.g., Baltic Sea)
- # Evaporation in Isolated Basins – evaporites
- # Hydrothermal Vents – brines high in salt
- # Precipitation & Dissolution – aragonite & calcite dissolution in deep ocean increase  $\text{Ca}^{2+}$  levels with precipitation elsewhere

# Exceptions to the Rule

(continued)

- # Anoxic Basins – bacterial reduction of  $\text{SO}_4^{2-}$  to  $\text{S}^{2-}$
- # Exchange at the Air-Sea interface – causes fractionation of many components
- # Freezing – sea ice can be deficient in one or more constituents causing local concentration anomalies
- # Interstitial Waters or Pore Waters – variety of processes many related to high surface areas in contact with water & anoxia

# Cl<sup>-</sup> has been Described as the Ultimate Conservative Tracer

- # Highest concentration in SW
- # Not biologically depleted
- # Not chemically limited
- # One of the longest Residence Times ( $1 \times 10^8$  yr)
- # Generally pretty boring
- # Oceanographers have used Cl<sup>-</sup> concentration to define the concentration of ocean water masses
- # Concept of Chlorinity = Cl<sup>-</sup> (+ Br<sup>-</sup>) content of SW

# Chlorinity (Cl)

- # Amount of  $\text{Cl}^-$ ,  $\text{Br}^-$  and  $\text{I}^-$  in grams, contained in 1 kg of seawater assuming  $\text{Br}^-$  and  $\text{I}^-$  replaced by  $\text{Cl}^-$
- # The number giving chlorinity in per mille of a seawater sample is by definition identical with the number giving the mass with unit gram of atomic weight silver just necessary to precipitate the halogens in 0.3285234 kg of the seawater sample (Jacobsen & Knudsen, 1940).

# Salinity (S)

- # Historical Definition - Total amount of solid material, in grams, contained in 1 kg of seawater when all carbonate has been converted to oxide, the bromide and iodine replaced by chlorine, and all organic matter completely oxidized
- # Practical Salinity Scale – Conductivity of seawater compared to KCl at 32.4356 g/kg (15 °C)

# Practical Salinity Scale (PSS 1978)

- #  $R_T = C(\text{sample})/C(\text{std seawater})$
- #  $C =$  conductivity at specified temp. & pressure
- # Formerly used units of parts per thousand ( $\text{‰}$ )
- # Unitless since based on a ratio
- # Often see PSU or practical salinity units
- # Calibrate instrumentation with SW standard

# Absolute Salinity ( $S_R$ )

## # SCOR/IAPSO

Scientific Committee on Oceanic Research

International Agency for the Physical Sciences of the Oceans

- WG 127 Thermodynamics & Equations of State of SW
- Density, Enthalpy, Entropy, Potential temp., Freezing temp.,
- Dissolved oxygen, Alkalinity,  $\text{TCO}_2$ , Ca, Silica

$$S_R = (35.16504 / 35) \text{ g/kg} \times S$$

# Precision in Salinity by Various Methods

- |  |             |
|--|-------------|
| 1) Composition Studies of major components | $\pm 0.01$  |
| 2) Evaporation to dryness                  | $\pm 0.01$  |
| 3) Chlorinity                              | $\pm 0.002$ |
| 4) Sound Speeds                            | $\pm 0.03$  |
| 5) Density                                 | $\pm 0.004$ |
| 6) Conductivity                            | $\pm 0.001$ |
| 7) Refractive index                        | $\pm 0.05$  |
| 8) Inductive Salinometer                   |             |

# Relationship between Salinity & Chlorinity

$$S = 1.80655 Cl$$

See Website for Salinity Handouts 1 - 4


# CTDs

Shown with  
optional cage,  
SBE 5T pump,  
& SBE 43 DO  
sensor

[www.seabird.com](http://www.seabird.com)


[www.valeport.co.uk](http://www.valeport.co.uk)





(a)

A vertical section showing the mean distribution of salinity in the western Atlantic Ocean, and two salinity depth profiles corresponding to locations A and B.



(b)

# Chemical Equilibria

General representation




Where uppercase letters are chemical species  
and lowercase letters are coefficients  
(i.e. # of atoms or moles)

# Equilibrium Constant

$$K = \frac{[C]^c [D]^d}{[A]^a [B]^b}$$

where [ ] = concentration, usually molar

# Solubility Equilibria



or by convention



# Solubility Product (equilibrium constant)

$$K_{sp} = \frac{[\text{Ba}^{2+}] [\text{SO}_4^{2-}]}{1} = [\text{Ba}^{2+}] [\text{SO}_4^{2-}]$$

$$K_{sp} = \frac{a_{\text{Ba}} a_{\text{SO}_4}}{1} = a_{\text{Ba}} a_{\text{SO}_4}$$

activity of solid is defined as = 1

# Solubility Calculated

Solubility (S) is the concentration of individual ions generated from an insoluble compound



$$S = [\text{Ba}^{2+}] = [\text{SO}_4^{2-}]$$

# Solubility Calculation (continued)

Given  $K_{SP} = [\text{Ba}^{2+}][\text{SO}_4^{2-}] = 2.0 \times 10^{-10}$

Then  $S = \sqrt{K_{SP}} = \sqrt{2.0 \times 10^{-10}} = 1.4 \times 10^{-5}$

So  $S = [\text{Ba}^{2+}] = [\text{SO}_4^{2-}] = 1.4 \times 10^{-5}$

# Activity Correction

$$K_{SP} = \frac{a_{Ba} a_{SO_4}}{1} = a_{Ba} a_{SO_4}$$

Since

$$a_{Ba} = \gamma_{Ba} [Ba^{2+}] \quad \& \quad a_{SO_4} = \gamma_{SO_4} [SO_4^{2-}]$$

Substituting

$$K_{SP} = a_{Ba} a_{SO_4} = \gamma_{Ba} [Ba^{2+}] \gamma_{SO_4} [SO_4^{2-}]$$

# Solubility Calculation (completed)

Since

$$K_{SP} = [\text{Ba}^{2+}][\text{SO}_4^{2-}] \quad \& \quad \gamma_{\text{Ba}} = \gamma_{\text{SO}_4}$$

Then

$$S = \sqrt{\frac{K_{SP}}{\gamma^2}}$$

To determine the solubility of  $\text{BaSO}_4$  in a solution containing other ions (like SW), you must calculate the activity coefficient ( $\gamma$ )