

Atmospheric Chemistry

- Structure and Circulation
- Gas Composition
- Photochemistry
- Case Study: Ozone






FIGURE 5.1. Changes in temperature in the layers of the atmosphere.




Thermosphere (Above 100 km)

N_2 and O_2 absorb UV light below 240 nm

Stratosphere (10 to 50 km)

O_3 absorbs UV light between 240 and 300 nm



Troposphere (below 10 km)

no UV light available to break O-O bonds
 > 300 nm of 120 kcal bond strength

Oxidizer is the OH^\cdot radical not O_3 or H_2O_2

FIGURE 5.1. Changes in temperature in the layers of the atmosphere.





FIGURE 5.3. Concentration of ozone in various layers of the atmosphere.

Surface Wind Bands



Adapted from Duxbury, Alyn C. and Alison B. Duxbury. *An Introduction to the World's Oceans*, 4/e.
Copyright © 1994 Wm. C. Brown Publishers, Dubuque, Iowa.





FIGURE 5.9

The movement of dust and particles in the northern hemisphere after the El Chichon Volcano eruption.

CH₄ CONCENTRATION (ppm)



CO CONCENTRATION (ppb)




FIGURE 5.8. Distribution of methane (top) and carbon monoxide (bottom) between the hemispheres.

Atmospheric Chemistry

- Structure and Circulation
- Gas Composition
- Photochemistry
- Case Study: Ozone

TABLE 5.1
Abundance of the Major
Conservative Atmospheric Gases^a

Mole fraction in dry air


Gas	(X_i)
N ₂	0.78084 ± 0.00004
O ₂	0.20946 ± 0.00002
Ar	$(9.34 \pm 0.01) \times 10^{-3}$
CO ₂	$(3.5 \pm 0.1) \times 10^{-4}$
Ne	$(1.818 \pm 0.004) \times 10^{-5}$
He	$(5.24 \pm 0.004) \times 10^{-6}$
Kr	$(1.14 \pm 0.01) \times 10^{-6}$
Xe	$(8.7 \pm 0.1) \times 10^{-8}$

^a Kester (1975).

TABLE 5.2**The Composition of Minor Gases in the Atmosphere**

Species	X_i actual	Reliability	Source	Sink
CH ₄	1.7×10^{-6}	High	Biog.	Photochem.
CO	$0.5\text{--}2 \times 10^{-7}$	Fair	Photo., anthr.	Photochem.
O ₃	5×10^{-8} (clean) 4×10^{-7} (polluted) 10^{-7} to 6×10^{-6} (stratosphere)	Fair	Photo	Photochem.
NO + NO ₂	$10^{-12}\text{--}10^{-8}$	Low	Lightn., anthr. photo.	Photochem.
HNO ₃	$10^{-11}\text{--}10^{-9}$	Low	Photo.	Rainout
NH ₃	$10^{-10}\text{--}10^{-9}$	Low	Biog.	Photo., rainout
N ₂ O	3×10^{-7}	High	Biog.	Photo.
H ₂	5×10^{-7}	High	Biog., photo.	Photo.
OH	$10^{-15}\text{--}10^{-12}$	Very low	Photo.	Photo.
HO ₂	$10^{-11}\text{--}10^{-13}$	Very low	Photo.	Photo.
H ₂ O ₂	$10^{-10}\text{--}10^{-18}$	Very low	Photo.	Rainout
H ₂ CO	$10^{-10}\text{--}10^{-9}$	Low	Photo.	Photo.
SO ₂	$10^{-11}\text{--}10^{-10}$	Fair	Anth., photo.,	Photo., volcanic
CS ₂	$10^{-11}\text{--}10^{-10}$	Low	Anthr., biol.,	Photo.
OCS	5×10^{-10}	Fair	Anthr., biol., photo.	Photo.
CH ₃ CCl ₃	$0.7\text{--}2 \times 10^{-10}$	Fair	Anthropogenic	Photo.




Atmospheric lifetimes of trace gases vary from a second to a century






Atmospheric Chemistry

- Structure and Circulation
- Gas Composition
- Photochemistry
- Case Study: Ozone

Production of OH Radicals





Removal Of OH Radicals





SUMMARY OF IMPORTANT NITROGEN OXIDE REACTIONS

(Species NO, NO₂, HNO₃)

Gas Phase





Aqueous Phase






Most important source of O₃ and other oxidants in the Troposphere.
Hydrocarbons are needed to produce HO₂.

SULFUR OXIDE CHEMISTRY

Gas



Aqueous



Atmospheric Chemistry

- Structure and Circulation
- Gas Composition
- Photochemistry
- Case Study: Ozone






FIGURE 5.37


The levels of ozone as a function of latitude.

Total Ozone for Aug 1, 1997



GSFC/918

Ozone Depletion Over Antarctica, Mean October Values at Halley Bay






FIGURE 5.34

The changes in the biologically damaging ultraviolet (UV) flux as a function of changes in ozone levels.

SUMMARY OF OZONE CHEMISTRY

FORMATION












PHOTO-DESTRUCTION





LOSS OF OZONE IN STRATOSPHERE


Chlorofluoro Carbons (CFC' s)



Scientific Support

F is found in Stratosphere

ClO appearance and O₃ loss in Antarctic








FIGURE 5.40

The relationship between O₃ and ClO in the Antarctic.


Annual Production of Fluorocarbons Reported to AFEAS (1980-2007)



CFCs and SF₆ in the Northern Hemisphere Atmosphere




Impact of Montreal Protocol on Chlorine Content of the Stratosphere



Without Controls 1996 Phaseout

Global Total Ozone Change

Changes from 1964-1980 average



Changes between 1980 and 2000

