


1. False. For example, take $f(x) = x^3$, then $f'(x) = 3x^2$ and $f'(0) = 0$, but $f(0) = 0$ is not a maximum or minimum; $(0, 0)$ is an inflection point.
2. False. For example, $f(x) = |x|$ has an absolute minimum at 0, but $f'(0)$ does not exist.
3. False. For example, $f(x) = x$ is continuous on $(0, 1)$ but attains neither a maximum nor a minimum value on $(0, 1)$. Don't confuse this with f being continuous on the *closed* interval $[a, b]$, which would make the statement true.
4. True. By the Mean Value Theorem, $f'(c) = \frac{f(1) - f(-1)}{1 - (-1)} = \frac{0}{2} = 0$. Note that $|c| < 1 \Leftrightarrow c \in (-1, 1)$.
5. True. This is an example of part (b) of the I/D Test.
6. False. For example, the curve $y = f(x) = 1$ has no inflection points but $f''(c) = 0$ for all c .
7. False. $f'(x) = g'(x) \Rightarrow f(x) = g(x) + C$. For example, if $f(x) = x + 2$ and $g(x) = x + 1$, then $f'(x) = g'(x) = 1$, but $f(x) \neq g(x)$.
8. False. Assume there is a function f such that $f(1) = -2$ and $f(3) = 0$. Then by the Mean Value Theorem there exists a number $c \in (1, 3)$ such that $f'(c) = \frac{f(3) - f(1)}{3 - 1} = \frac{0 - (-2)}{2} = 1$. But $f'(x) > 1$ for all x , a contradiction.
9. True. The graph of one such function is sketched.



10. False. At any point $(a, f(a))$, we know that $f'(a) < 0$. So since the tangent line at $(a, f(a))$ is not horizontal, it must cross the x -axis—at $x = b$, say. But since $f''(x) > 0$ for all x , the graph of f must lie above all of its tangents; in particular, $f(b) > 0$. But this is a contradiction, since we are given that $f(x) < 0$ for all x .
11. True. Let $x_1 < x_2$ where $x_1, x_2 \in I$. Then $f(x_1) < f(x_2)$ and $g(x_1) < g(x_2)$ [since f and g are increasing on I], so $(f + g)(x_1) = f(x_1) + g(x_1) < f(x_2) + g(x_2) = (f + g)(x_2)$.
12. False. $f(x) = x$ and $g(x) = 2x$ are both increasing on $(0, 1)$, but $f(x) - g(x) = -x$ is not increasing on $(0, 1)$.
13. False. Take $f(x) = x$ and $g(x) = x - 1$. Then both f and g are increasing on $(0, 1)$. But $f(x)g(x) = x(x - 1)$ is not increasing on $(0, 1)$.

14. True. Let $x_1 < x_2$ where $x_1, x_2 \in I$. Then $0 < f(x_1) < f(x_2)$ and $0 < g(x_1) < g(x_2)$ [since f and g are both positive and increasing]. Hence, $f(x_1)g(x_1) < f(x_2)g(x_1) < f(x_2)g(x_2)$. So fg is increasing on I .
15. True. Let $x_1, x_2 \in I$ and $x_1 < x_2$. Then $f(x_1) < f(x_2)$ [f is increasing] $\Rightarrow \frac{1}{f(x_1)} > \frac{1}{f(x_2)}$ [f is positive] $\Rightarrow g(x_1) > g(x_2) \Rightarrow g(x) = 1/f(x)$ is decreasing on I .
16. False. If f is even, then $f(x) = f(-x)$. Using the Chain Rule to differentiate this equation, we get $f'(x) = f'(-x) \frac{d}{dx}(-x) = -f'(-x)$. Thus, $f'(-x) = -f'(x)$, so f' is odd.
17. True. If f is periodic, then there is a number p such that $f(x+p) = f(x)$ for all x . Differentiating gives $f'(x) = f'(x+p) \cdot (x+p)' = f'(x+p) \cdot 1 = f'(x+p)$, so f' is periodic.
18. False. The most general antiderivative of $f(x) = x^{-2}$ is $F(x) = -1/x + C_1$ for $x < 0$ and $F(x) = -1/x + C_2$ for $x > 0$ [see Example 1 in Section 4.7].
19. True. By the Mean Value Theorem, there exists a number c in $(0, 1)$ such that $f(1) - f(0) = f'(c)(1 - 0) = f'(c)$. Since $f'(c)$ is nonzero, $f(1) - f(0) \neq 0$, so $f(1) \neq f(0)$.