

Bipolar I disorder

- A. Criteria have been met for at least one manic episode (Criteria A-D under “Manic Episode” above) (LMS Unit 7.1b).
- B. The occurrence of the manic and major depressive episode(s) is not better explained by schizoaffective disorder, schizophrenia, schizophreniform disorder, delusional disorder, or other specified or unspecified schizophrenia spectrum and other psychotic disorder.

Specify:

With anxious distress

With mixed features

With rapid cycling

With melancholic features

With atypical features

With mood-congruent psychotic features

With mood-incongruent psychotic features

With catatonia

With peripartum onset

With seasonal pattern

Bipolar II disorder

- A. Criteria have been met for at least one hypomanic episode (Criteria A-F under “Hypomanic Episode” above) (LMS 7.1c) and at least one major depressive episode (Criteria A-C under “Major Depressive Episode” above) (LMS 7.1a).
- B. There has never been a manic episode.
- C. The occurrence of this manic and major depressive episode(s) is not better explained by schizoaffective disorder, schizophrenia, schizophreniform disorder, delusional disorder, or other specified or unspecified schizophrenia spectrum and other psychotic disorder.
- D. The symptoms of depression or the unpredictability caused by frequent alternation between periods of depression and hypomania causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.

Specify:

With anxious distress

With mixed features

With rapid cycling

With mood-congruent psychotic features

With mood-incongruent psychotic features

With catatonia

With peripartum onset

With seasonal pattern

