

## Erectile disorder

A. At least one of the three following symptoms must be experienced on almost all (approximately 75%-100%) occasions of sexual activity (in identified situational contexts or, if generalized, in all contexts):

1. Marked difficulty in obtaining an erection during sexual activity.
2. Marked difficulty in maintaining an erection until the completion of sexual activity
3. Marked decrease in erectile rigidity.

B. The symptoms in Criterion A have persisted for a minimum duration of approximately 6 months.

C. The symptoms in Criterion A cause clinically significant distress in the individual.

D. The sexual dysfunction is not better explained by a nonsexual mental disorder or as a consequence of severe relationship distress or other significant stressors and is not attributable to the effects of a substance/medication or another medical condition.

*Specify whether:*

**Lifelong:** The disturbance has been present since the individual became sexually active.

**Acquired:** The disturbance began after a period of relatively normal sexual function.

*Specify whether:*

**Generalized:** Not limited to certain types of stimulation, situations, or partners.

**Situational:** Only occurs with certain types of stimulation, situations, or partners.

*Specify whether:*

**Mild:** Evidence of mild distress over the symptoms in Criterion A

**Moderate:** Evidence of moderate distress over the symptoms in Criterion A

**Severe:** Evidence of severe or extreme distress over the symptoms in Criterion A