

Female orgasmic disorder

- A. Presence of either of the following symptoms and experienced on almost all or all (approximately 75%-100%) occasions of sexual activity (in identified situational contexts or, if generalized, in all contexts):
 - 1. Marked delay in, marked infrequency of, or absence of orgasm.
 - 2. Markedly reduced intensity of orgasmic sensations.
- B. The symptoms in Criterion A have persisted for a minimum duration of approximately 6 months.
- C. The symptoms in Criterion A cause clinically significant distress in the individual.

The sexual dysfunction is not better explained by a nonsexual mental disorder or as a consequence of severe relationship distress (e.g., partner violence) or other significant stressors and is not attributable to the effects of a substance/medication or another medical condition.

Specify whether:

Lifelong: The disturbance has been present since the individual became sexually active.

Acquired: The disturbance began after a period of relatively normal sexual function.

Specify whether:

Generalized: Not limited to certain types of stimulation, situations, or partners.

Situational: Only occurs with certain types of stimulation, situations, or partners.

Specify if:

Never experienced an orgasm under any situation.

Specify whether:

Mild: Evidence of mild distress over the symptoms in Criterion A

Moderate: Evidence of moderate distress over the symptoms in Criterion A

Severe: Evidence of severe or extreme distress over the symptoms in Criterion A