Panic disorder

A. Recurrent unexpected panic attacks. A panic attack is an abrupt surge of intense fear or intense discomfort that reaches a peak within minutes, and during which time four (or more) of the following symptoms occur:

Note: The abrupt surge can occur from a calm state or an anxious state.

- 1. Palpitations, pounding heart, or accelerated heart rate.
- 2. Sweating.
- 3. Trembling or shaking.
- 4. Sensations of shortness of breath or smothering.
- 5. Feelings of choking.
- 6. Chest pain or discomfort.
- 7. Nausea or abdominal distress.
- 8. Feeling dizzy, unsteady, light-headed, or faint.
- 9. Chills or heat sensations.
- 10. Paresthesias (numbness or tingling sensations).
- 11. Derealization (feelings of unreality) or depersonalization (being detached from oneself).
- 12. Fear of losing control or "going crazy."
- 13. Fear of dying.

Note: Culture-specific symptoms (e.g., tinnitus, neck soreness, headache, uncontrollable screaming or crying) may be seen. Such symptoms should not count as one of the four required symptoms.

- B. At least one of the attacks has been followed by 1 month (or more) of one or both of the following:
 - 1. Persistent concern or worry about additional panic attacks or their consequences (e.g., losing control, having a heart attack, "going crazy").
 - 2. A significant maladaptive change in behavior related to the attacks (e.g., behaviors designed to avoid having panic attacks, such as avoidance of exercise or unfamiliar situations).
- C. The disturbance is not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication) or another medical condition (e.g., hyperthyroidism, cardiopulmonary disorders).
- D. The disturbance is not better explained by another mental disorder (e.g., the panic attacks do not occur only in response to feared social situations, as In social anxiety disorder; in response to circumscribed phobic objects or situations, as in specific phobia; in response to obsessions, as in obsessive-compulsive disorder; in response to reminders of traumatic events, as in posttraumatic stress disorder; or in response to separation from attachment figures, as in separation anxiety disorder).